

Proyecciones Journal of Mathematics  
Vol. 35, N° 3, pp. 235-244, September 2016.  
Universidad Católica del Norte  
Antofagasta - Chile  
DOI: 10.4067/S0716-09172016000300001

## An alternative proof of a Tauberian theorem for Abel summability method

*İbrahim Çanak*  
*Ege University, Turkey*  
*and*

*Ümit Totur*  
*Adnan Menderes University, Turkey*  
*Received : December 2014. Accepted : July 2016*

### Abstract

*Using a corollary to Karamata's main theorem [Math. Z. 32 (1930), 319–320], we prove that if a slowly decreasing sequence of real numbers is Abel summable, then it is convergent in the ordinary sense.*

**Subjclass [2010] :** 40A05; 40E05; 40G10.

**Keywords :** Abel summability, slowly decreasing sequences, Tauberian conditions and theorems.

## 1. Introduction

A number of authors such as Schmidt [9], Maddox [6], Móricz [8], and Talo and Başar [11] have proved several Tauberian theorems for some summability methods for which slowly decreasing condition for sequences is a Tauberian condition. Schmidt [9] obtained that the slowly decreasing condition for sequences of real numbers is a Tauberian condition for Abel summability. Maddox [6] introduced the slowly decreasing sequence in an ordered linear space and proved that a Cesàro summable sequence is convergent if it is slowly decreasing in an ordered linear space. Móricz [8] established a Tauberian theorem which states that ordinary convergence of a sequence follows from its statistical Cesàro summability if it is slowly decreasing. Talo and Başar [11] introduced the concept of slowly decreasing sequences for fuzzy numbers and they proved that the slowly decreasing condition for sequences is a Tauberian condition for the statistical convergence and Cesàro summability for sequences of fuzzy numbers.

Littlewood [5] proved that  $n(u_n - u_{n-1}) = O(1)$  is a Tauberian condition for Abel summability of  $(u_n)$ . But his proof was complicated and based on the repeated differentiation. A first clever and surprisingly simple proof based on Weierstrass approximation theorem of Littlewood's theorem was given by Karamata [2].

The main purpose of this study is to give an alternative simpler proof of the following Tauberian theorem which is more general than Littlewood's theorem [5] for Abel summability method.

**Theorem 1.1.** *If  $(u_n)$  is Abel summable to  $s$  and slowly decreasing, then  $\lim_n u_n = s$ .*

To prove Theorem 1.1, we first obtain Cesàro convergence of the generator sequence of a given sequence  $(u_n)$  by means of a corollary to Karamata's main Theorem, and then recover convergence of  $(u_n)$  by Tauber's second theorem [12].

Our proof is much easier than the existing one and uses the well known results in Tauberian theory. For a different proof of Theorem 1.1, see [1].

## 2. Preliminaries

For a sequence  $u = (u_n)$  of real numbers, we write  $(u_n)$  in terms of  $(v_n)$  as

$$(2.1) \quad u_n = v_n + \sum_{k=1}^n \frac{v_k}{k} + u_0, \quad (n = 1, 2, \dots)$$

where  $v_n = \frac{1}{n+1} \sum_{k=1}^n k(u_k - u_{k-1})$ . The sequence  $(v_n)$  is called a generator sequence of  $(u_n)$ . We note that  $\sigma_n^{(1)}(u) = \frac{1}{n+1} \sum_{k=0}^n u_k = u_0 + \sum_{k=1}^n \frac{v_k}{k}$ .

Let  $u = (u_n)$  be a sequence of real numbers. For each nonnegative integer  $m$ , we define  $\sigma_n^{(m)}(u)$  by

$$\sigma_n^{(m)}(u) = \begin{cases} \frac{1}{n+1} \sum_{k=0}^n \sigma_k^{(m-1)}(u) & , m \geq 1 \\ u_n & , m = 0 \end{cases}$$

A sequence  $(u_n)$  is said to be Abel summable to  $s$  if  $u_0 + \sum_{n=1}^{\infty} (u_n - u_{n-1})x^n$  converges for  $0 < x < 1$ , and tends to  $s$  as  $x \rightarrow 1^-$ .

A sequence  $(u_n)$  is called  $(A, m)$  summable to  $s$  if  $(\sigma_n^{(m)}(u))$  is Abel summable to  $s$ . If  $m = 0$ , then  $(A, m)$  summability reduces to Abel summability. It is clear that Abel summability of  $(u_n)$  implies  $(A, m)$  summability of  $(u_n)$ .

Throughout this work, the symbol  $[\lambda n]$  denotes the integral part of the product  $\lambda n$ .

A sequence  $(u_n)$  is said to be slowly decreasing [9] if

$$(2.2) \quad \lim_{\lambda \rightarrow 1^+} \liminf_{n \rightarrow \infty} \min_{n+1 \leq k \leq [\lambda n]} (u_k - u_n) \geq 0$$

or equivalently [8],

$$(2.3) \quad \lim_{\lambda \rightarrow 1^-} \liminf_{n \rightarrow \infty} \min_{[\lambda n] + 1 \leq k \leq n} (u_n - u_k) \geq 0.$$

Notice that  $(u_n)$  is slowly decreasing if the classical one-sided Tauberian condition of Landau [1] is satisfied, that is, there exists a positive constant  $C > 0$  such that

$$(2.4) \quad n(u_n - u_{n-1}) \geq -C$$

for all nonnegative  $n$ . Indeed, for any  $k > n$ , we have

$$u_k - u_n = \sum_{j=n+1}^k (u_j - u_{j-1}) \geq -C \sum_{j=n+1}^k \frac{1}{j} \geq -C \log \left( \frac{k}{n} \right)$$

whence we conclude that

$$\liminf_{n \rightarrow \infty} \min_{n+1 \leq k \leq [\lambda n]} (u_k - u_n) \geq -C \log \lambda, \quad \lambda > 1.$$

Taking  $\lambda \rightarrow 1^+$ , we have the inequality (2.2).

Note that we used  $C$  to denote a constant, possibly different at each occurrence.

A sequence  $(u_n)$  is slowly increasing if and only if  $(-u_n)$  is slowly decreasing, and an equivalent definition of a slowly increasing sequence as follows:

A sequence  $(u_n)$  is said to be slowly increasing if

$$(2.5) \quad \lim_{\lambda \rightarrow 1^+} \limsup_{n \rightarrow \infty} \max_{n+1 \leq k \leq [\lambda n]} (u_k - u_n) \leq 0.$$

The condition (2.5) is reformulated as follows (see [8]):

$$(2.6) \quad \lim_{\lambda \rightarrow 1^-} \limsup_{n \rightarrow \infty} \max_{[\lambda n]+1 \leq k \leq n} (u_n - u_k) \leq 0.$$

It is plain that a sequence  $(u_n)$  is said to be slowly oscillating if and only if  $(u_n)$  is both slowly increasing and slowly decreasing. Notice that each of the conditions (2.2) and (2.5) is necessary for convergence (see [4]).

If  $(u_n)$  converges to  $s$ , then  $(u_n)$  is Abel summable to  $s$ . However, the converse of this statement is not always true. Note that Abel summability of  $(u_n)$  implies convergence of  $(u_n)$  under certain additional hypotheses called Tauberian conditions. Any theorem which states that convergence of sequence  $(u_n)$  follows from Abel summability of  $(u_n)$  and some Tauberian condition(s) is called a Tauberian theorem for Abel summability method.

### 3. Corollary to Karamata's Main Theorem and Lemmas

Our proof is based on the following corollary to Karamata's main theorem and three Lemmas.

**Corollary to Karamata's Main Theorem.** ([2]) If  $u = (u_n)$  is Abel summable to  $s$  and  $u_n \geq -C$  for some nonnegative  $C$ , then  $\lim_n \sigma_n^{(1)}(u) = s$ .

**Lemma 3.1.** ([3]) If, for  $x \rightarrow 1^-$ , a function  $f(x)$ , which is integrable in  $[0, 1]$ , satisfies the limiting relation

$$(3.1) \quad (1-x)^2 f(x) \rightarrow s,$$

then, for  $x \rightarrow 1^-$ , we also have

$$(3.2) \quad (1-x) \int_0^x f(t) dt \rightarrow s.$$

The next lemma gives a necessary condition for a slowly decreasing sequence in terms of the generator sequence  $(v_n)$ .

**Lemma 3.2.** ([7]) *If  $(u_n)$  is slowly decreasing, then  $v_n \geq -C$  for some  $C$ , where  $v_n = \frac{1}{n+1} \sum_{k=1}^n k(u_k - u_{k-1})$ .*

Next, we represent the difference  $u_n - \sigma_n^{(1)}(u)$  in two different ways.

**Lemma 3.3.** ([10]) *Let  $u = (u_n)$  be a sequence of real numbers.*

(i) *For  $\lambda > 1$  and sufficiently large  $n$ ,*

$$(3.3) \quad u_n - \sigma_n^{(1)}(u) = \frac{[\lambda n] + 1}{[\lambda n] - n} \left( \sigma_{[\lambda n]}^{(1)}(u) - \sigma_n^{(1)}(u) \right) - \frac{1}{[\lambda n] - n} \sum_{k=n+1}^{[\lambda n]} (u_k - u_n).$$

(ii) *For  $0 < \lambda < 1$  and sufficiently large  $n$ ,*

$$(3.4) \quad u_n - \sigma_n^{(1)}(u) = \frac{[\lambda n] + 1}{n - [\lambda n]} \left( \sigma_n^{(1)}(u) - \sigma_{[\lambda n]}^{(1)}(u) \right) + \frac{1}{n - [\lambda n]} \sum_{k=[\lambda n]+1}^n (u_n - u_k).$$

#### 4. Proof of Theorem 1.1

**Proof.** Since  $(u_n)$  is Abel summable to  $s$ , then  $(\sigma_n^{(1)}(u))$  is also Abel summable to  $s$ . Hence, we conclude by (2.1) that  $(v_n) = \left( \frac{1}{n+1} \sum_{k=0}^n k(u_k - u_{k-1}) \right)$  is Abel summable to zero by Lemma 3.1. It follows by Lemma 3.2 that there exists a nonnegative  $C$  such that

$$(4.1) \quad v_n \geq -C.$$

Taking (4.1) and the fact that  $(v_n)$  is Abel summable to zero into account, we obtain by Corollary to Karamata's Main Theorem that  $\sigma_n^{(1)}(v) =$

$o(1)$  as  $n \rightarrow \infty$ . Since  $(\sigma_n^{(1)}(u))$  is Abel summable to  $s$  and  $\sigma_n^{(1)}(v) = o(1)$  as  $n \rightarrow \infty$ , we have that  $(\sigma_n^{(1)}(u))$  converges to  $s$  by Tauber's second theorem [12].

By the fact that every convergent sequence is slowly increasing, we have

$(\sigma_n^{(1)}(u))$  is slowly increasing. Thus,  $(-\sigma_n^{(1)}(u))$  is slowly decreasing. Since  $(s_n)$  is slowly decreasing,  $(v_n)$  is slowly decreasing.

By Lemma 3.3 (i), we have

$$v_n - \sigma_n^{(1)}(v) = \frac{[\lambda n] + 1}{[\lambda n] - n} \left( \sigma_{[\lambda n]}^{(1)}(v) - \sigma_n^{(1)}(v) \right) - \frac{1}{[\lambda n] - n} \sum_{k=n+1}^{[\lambda n]} (v_k - v_n). \quad (4.2)$$

It is easy to verify that for  $\lambda > 1$  and sufficiently large  $n$ ,

$$\frac{\lambda}{2(\lambda - 1)} \leq \frac{[\lambda n] + 1}{[\lambda n] - n} \leq \frac{3\lambda}{2(\lambda - 1)}. \quad (4.3)$$

By  $\sigma_n^{(1)}(v) = o(1)$  as  $n \rightarrow \infty$  and (4.10), for all  $\lambda > 1$ ,

$$\lim_{n \rightarrow \infty} \frac{[\lambda n] + 1}{[\lambda n] - n} \left( \sigma_{[\lambda n]}^{(1)}(v) - \sigma_n^{(1)}(v) \right) = 0. \quad (4.4)$$

By (4.2) and (4.3), we have

$$v_n - \sigma_n^{(1)}(v) \leq \frac{[\lambda n] + 1}{[\lambda n] - n} \left( \sigma_{[\lambda n]}^{(1)}(v) - \sigma_n^{(1)}(v) \right) - \min_{n+1 \leq k \leq [\lambda n]} (v_k - v_n). \quad (4.5)$$

Taking  $\limsup$  of both sides of (4.5), we have

$$\begin{aligned} \limsup_n (v_n - \sigma_n^{(1)}(v)) &\leq \limsup_{n \rightarrow \infty} \frac{[\lambda n] + 1}{[\lambda n] - n} \left( \sigma_{[\lambda n]}^{(1)}(v) - \sigma_n^{(1)}(v) \right) \\ &\quad - \liminf_n \min_{n+1 \leq k \leq [\lambda n]} (v_k - v_n). \end{aligned} \quad (4.6)$$

The inequality (4.6) becomes

$$(4.7) \quad \limsup_n (v_n - \sigma_n^{(1)}(v)) \leq -\liminf_n \min_{n+1 \leq k \leq [\lambda n]} (v_k - v_n)$$

by (4.4). Taking  $\lambda \rightarrow 1^+$  in (4.7), we have

$$(4.8) \quad \limsup_n (v_n - \sigma_n^{(1)}(v)) \leq 0$$

by (2.2).

By Lemma 3.3 (ii), we have

$$(4.9) \quad v_n - \sigma_n^{(1)}(v) = \frac{[\lambda n] + 1}{n - [\lambda n]} \left( \sigma_n^{(1)}(v) - \sigma_{[\lambda n]}^{(1)}(v) \right) + \frac{1}{n - [\lambda n]} \sum_{k=[\lambda n]+1}^n (v_n - v_k).$$

It is easy to verify that for  $0 < \lambda < 1$  and sufficiently large  $n$ ,

$$(4.10) \quad \frac{\lambda}{2(1-\lambda)} \leq \frac{[\lambda n] + 1}{n - [\lambda n]} \leq \frac{3\lambda}{2(1-\lambda)}.$$

By  $\sigma_n^{(1)}(v) = o(1)$  as  $n \rightarrow \infty$  and (4.10), for all  $0 < \lambda < 1$ ,

$$(4.11) \quad \lim_{n \rightarrow \infty} \frac{[\lambda n] + 1}{n - [\lambda n]} \left( \sigma_n^{(1)}(v) - \sigma_{[\lambda n]}^{(1)}(v) \right) = 0.$$

By (4.9) and (4.10), we have

$$(4.12) \quad v_n - \sigma_n^{(1)}(v) \geq \frac{[\lambda n] + 1}{n - [\lambda n]} \left( \sigma_n^{(1)}(v) - \sigma_{[\lambda n]}^{(1)}(v) \right) + \min_{[\lambda n]+1 \leq k \leq n} (v_n - v_k).$$

Taking  $\liminf$  of both sides of (4.12), we have

$$(4.13) \quad \begin{aligned} \liminf_n (v_n - \sigma_n^{(1)}(v)) &\geq \liminf_n \frac{[\lambda n] + 1}{n - [\lambda n]} \left( \sigma_n^{(1)}(v) - \sigma_{[\lambda n]}^{(1)}(v) \right) \\ &+ \liminf_n \min_{[\lambda n]+1 \leq k \leq n} (v_n - v_k). \end{aligned}$$

The inequality (4.13) becomes

$$(4.14) \quad \liminf_n (v_n - \sigma_n^{(1)}(v)) \geq \liminf_n \min_{[\lambda n]+1 \leq k \leq n} (v_n - v_k)$$

by (4.11).

Taking  $\lambda \rightarrow 1^-$  in (4.14), we have

$$(4.15) \quad \liminf_n (v_n - \sigma_n^{(1)}(v)) \geq 0$$

by (2.3).

Combining (4.8) and (4.15) yields that  $v_n = o(1)$  as  $n \rightarrow \infty$ . Since  $(u_n)$  is Abel summable to  $s$  and  $v_n = o(1)$  as  $n \rightarrow \infty$ ,  $\lim_n u_n = s$  by Tauber's second theorem [12]. This completes the proof.  $\square$

Using Theorem 1.1, we show that slow decrease of  $(u_n)$  is also a Tauberian condition for  $(A, m)$  summability method.

**Theorem 4.1.** *If  $(u_n)$  is  $(A, m)$  summable to  $s$  and slowly decreasing, then  $\lim_n u_n = s$ .*

**Proof.** Let  $(u_n)$  be slowly decreasing. Then, we have  $v_n \geq -C$  for some  $C$  by Lemma 3.2. Since  $n(\sigma_n^{(1)}(u) - \sigma_{n-1}^{(1)}) = v_n$  for all nonnegative  $n$ , we conclude that  $(\sigma_n^{(1)}(u))$  is slowly decreasing if we replace  $u_n$  in (2.4) by  $\sigma_n^{(1)}(u)$ .

It easily follows that  $(\sigma_n^{(m)}(u))$  is slowly decreasing for each nonnegative  $m$ .

Since  $(u_n)$  is  $(A, m)$  summable to  $s$ , we have

$$(4.16) \quad \lim_n \sigma_n^{(m)}(u) = s$$

by Theorem 1.1. By definition, we have

$$(4.17) \quad \sigma_n^{(m)}(u) = \sigma_n^{(1)}(\sigma^{(m-1)}(u)).$$

From (4.16) and (4.17) it follows that  $(u_n)$  is  $(A, m-1)$  summable to  $s$ . Since  $(\sigma_n^{(m-1)}(u))$  is slowly decreasing, we have  $\lim_n \sigma_n^{(m-1)}(u) = s$  by Theorem 1.1. Continuing in this way, we obtain that  $\lim_n u_n = s$ .  $\square$



## References

- [1] G. H. Hardy, Divergent series, Oxford University Press, (1948).
- [2] J. Karamata, Über die Hardy-Littlewoodschen Umkehrungen des Abelschen Stetigkeitssatzes, Math. Z., 32, pp. 319–320, (1930).
- [3] K. Knopp, Theory and application of infinite series, Dover Publications, (1990).
- [4] J. Korevaar, Tauberian theory, Springer, 2004.
- [5] J. E. Littlewood, The converse of Abel's theorem on power series, London M. S. Proc. 2 (9), pp. 434–448, (1911).
- [6] I. J. Maddox, A Tauberian theorem for ordered spaces, Analysis, 9, (3), pp. 297–302, (1989).
- [7] G. A. Mikhalin, Theorem of Tauberian type for  $(J, p_n)$  summation methods, Ukrain. Mat. Zh. 29 (1977), 763–770. English translation: Ukrain. Math. J. 29, pp. 564–569, (1977).
- [8] F. Móricz, Ordinary convergence follows from statistical summability  $(C, 1)$  in the case of slowly decreasing or oscillating sequences, Colloq. Math. 99, (2), pp. 207–219, (2004).
- [9] R. Schmidt, Über divergente Folgen und lineare Mittelbildungen, Math. Z. 22 (1), pp. 89–152, (1925).
- [10] Č. V. Stanojević, V. B. Stanojević, Tauberian retrieval theory, Publ. Inst. Math. (Beograd) (N.S.) 71 (85), pp. 105–111, (2002).
- [11] Ö. Talo, F. Başar, On the slowly decreasing sequences of fuzzy numbers, Abstr. Appl. Anal. Art. ID 891986, 7, pp. ..., (2013).
- [12] A. Tauber, Ein satz aus der theorie der unendlichen reihen, Monatsh. f. Math. u. Phys. 7, pp. 273–277, (1897).

**İbrahim Çanak**

Department of Mathematics

Ege University

Izmir

Turkey

e-mail : [ibrahim.canak@ege.edu.tr](mailto:ibrahim.canak@ege.edu.tr)

and

**Ümit Totur**

Department of Mathematics

Adnan Menderes University

Aydin

Turkey

e-mail : [utotur@adu.edu.tr](mailto:utotur@adu.edu.tr)