

Stability and boundedness in differential systems of third order with variable delay

Cemil Tunc

Yuzuncu University, Turkey

Received : April 2016. Accepted : June 2016

Abstract

In this paper, we consider a non-linear system of differential equations of third order with variable delay. We discuss the globally asymptotic stability/uniformly stability, boundedness and uniformly boundedness of solutions for the considered system. The technique of proofs involves defining an appropriate Lyapunov functional. The obtained results include and improve the results in literature.

Keywords : *Globally asymptotic stability, boundedness, Lyapunov functional, delay, differential system, third order.*

2010 Mathematics Subject Classifications : *34K12, 34K20.*

1. Introduction

In the last years, there is a good amount of interest in the qualitative behaviors of ordinary and functional differential equations of third order without and with delay, see the book of Reissig et al. [1] as a good survey for the works done by 1974 and the papers of Ademola and Arawomo ([2], [3], [4],[5]), Ademola et al. [6], Afuwape and Castellanos [7], Afuwape and Omeike [8], Ahmad and Rama Mohana Rao [9], Bai and Guo [10], Ezeilo [11], Ezeilo and Tejumola [12], Graef et al. ([13], [14]), Graef and Tunc [15], Korkmaz and Tunc [16], Mahmoud and Tunc [17], Ogundare [18], Ogundare et al. [19], Olutimo [20], Omeike [21], Qian [22], Remili and Oudjedi [23], Sadek [24], Swick [25], Tejumola and Tchegnani [26], Tunc ([27]-[44]), Tunc and Ates [45], Tunc and Gozen [46], Tunc and Mohammed [47], Tunc and Tunc [48], Zhang and Yu [49], Zhu [50] and theirs references. However, to the best of our knowledge from the literature, by this time, no attention was given to the investigation of the globally asymptotic stability/uniformly stability, boundedness and uniformly boundedness in the systems of nonlinear functional differential equations of third order with variable delay, except the recent work of Omeike [21].

Besides, it is well known that differential equations of third order play extremely important and useful roles in many scientific areas such as atomic energy, biology, chemistry, control theory, economy, engineering, information theory, mechanics, medicine, physics, etc.. Indeed, we can find applications such as nonlinear oscillations in Afuwape et al. [51], Andres [52], Fridedrichs [53], physical applications in Animalu and Ezeilo [54], non-resonant oscillations in Ezeilo and Onyia [55], prototypical examples of complex dynamical systems in a high-dimensional phase space, displacement in a mechanical system, velocity, acceleration in Chlouverakis and Sprott [56], Eichhorn et al. [57], Linz [58], the biological model and other models in Cronin-Scanlon [59], problems in biomathematics in Chlouverakis and Sprott [56], electronic theory in Rauch [60], and etc. Further, we refer the readers to the book of Smith [61] for some important applications of delay differential equation in sciences, biomathematics, engineering, and etc..

In 2015, Omeike [21] investigated the stability and boundedness of nonlinear differential system of third order with variable delay, $\tau(t)$:

$$(1.1) \quad X''' + AX'' + BX' + H(X(t - \tau(t))) = P(t).$$

In this paper, we consider nonlinear differential system of the third

order with variable delay, $\tau(t)$:

$$X''' + AX'' + G(X'(t - \tau(t))) + H(X(t - \tau(t))) = F(t, X, X', X''), \quad (1.2)$$

where $X \in \mathcal{R}^n$, $t \in [0, \infty)$, $\mathcal{R}^+ = [0, \infty)$, $\tau(t)$ is a continuous differentiable function such that $0 \leq \tau(t) \leq \tau_0$ is a positive constant and $\tau'(t) \leq \tau_1$, ($0 < \tau_1 < 1$), A is an $n \times n$ - constant symmetric matrix, $G, H : \mathcal{R}^n \rightarrow \mathcal{R}^n$ are continuous differentiable functions with $G(0) = H(0) = 0$ such that the Jacobian matrices $J_G(X')$ and $J_H(X)$ exist and are symmetric and continuous, that is,

$$J_G(X') = \left(\frac{\partial g_i}{\partial x'_j} \right), J_H(X) = \left(\frac{\partial h_i}{\partial x_j} \right), (i, j = 1, 2, \dots, n),$$

exist and are symmetric and continuous, where $(x_1, x_2, \dots, x_n)$, $(x'_1, x'_2, \dots, x'_n)$, (g_i) and (h_i) are components of X , X' , G and H , respectively, $F : \mathcal{R}^+ \times \mathcal{R}^n \times \mathcal{R}^n \times \mathcal{R}^n \rightarrow \mathcal{R}^n$ is a continuous function and the primes in Eq. (1.2) indicate differentiation with respect to t , $t \geq t_0 \geq 0$.

It is more convenient to consider not Eq. (1.2) itself, but rather the system

$$\begin{aligned} X'_1 &= X_2, X'_2 = X_3, \\ X'_3 &= -AX_3 - G(X_2) - H(X_1) + \int_{t-\tau(t)}^t J_G(X_2(s))X_3(s)ds \\ (1.3) \quad &+ \int_{t-\tau(t)}^t J_H(X_1(s))X_2(s)ds + F(t, X_1, X_2, X_3). \end{aligned}$$

System (1.3) is obtained from Eq. (1.2) by setting $X = X_1$, $X' = X_2$ and $X'' = X_3$.

The continuity of the functions τ, H, G and F is a sufficient condition to guarantee the existence of solutions of Eq. (1.2). Besides, we assume that the functions H, G and F satisfy a Lipschitz condition with respect to their respective arguments, like X, X' and X'' . In this case, the uniqueness of solutions of Eq. (1.2) is guaranteed.

The motivation of this paper comes from the results established by Omeike [21], the mentioned books, papers and their references. The main

purpose of this paper is to get new the globally asymptotic stability /uniformly stability, boundedness and uniformly boundedness results in Eq. (1.2) by defining a suitable new Lyapunov functional. By this paper, we extend and improve the stability and boundedness results of Omeike [21], and we give additional two results to that in Omeike [21], like uniformly stability and uniformly boundedness results. It follows that if we choose $G(X'(t - \tau(t))) = BX'(t)$, B is an $n \times n$ -constant symmetric matrix, and $F(t, X, X', X'') = P(t)$ in Eq. (1.2), then Eq. (1.2) reduces to Eq. (1.1), which is discussed by Omeike [21]. This means that instead of the linear term $BX'(t)$ in Eq. (1.1), we take the non-linear term $G(X'(t - \tau(t)))$ which includes a variable delay $\tau(t)$, and we also take a nonlinear generalization of the term $P(t)$ in Eq. (1.1) like $F(t, X, X', X'')$. Probably, these cases seem as similarity of Eq. (1.1) and Eq. (1.2).

However, till now, throughout all the papers published in the literature, no author discussed the stability and boundedness of the solutions when we take the possible second term in Eq. (1.1), $G(X')$ or $BX'(t)$ as a non-linear term with a deviating argument like $G(X'(t - \tau(t)))$. To the best of our knowledge, it is not easy to discuss the topic for Eq. (1.2). The possible reason is that the construction or definition of a suitable Lyapunov function or functional for higher differential systems remains as an open problem in the literature by this time. This case is more difficult for the functional differential systems of higher order with variable delay. The choice of the second term in Eq.(1.2) as $G(X'(t - \tau(t)))$ is important and the discussion of the problems for this case is some hard. This means that, in view of the whole mentioned discussion, it is worth to discuss the globally asymptotic stability /uniformly stability, boundedness and uniformly boundedness to Eq. (1.2).

Besides, this paper may be useful for researchers working on the qualitative behaviors of solutions of third differential equations and completes that in the literature. These cases show the novelty and originality of the present paper.

Consider general delay differential system

$$\dot{x} = f(x_t), x_t = x(t + \theta), -r \leq \theta < 0, t \geq 0.$$

Let $C = C([-r, 0], \mathcal{R}^n)$ denote the space of continuous function from $[-r, 0]$ into $\mathcal{R}^n$ and assume that $f : C \rightarrow \mathcal{R}^n$ is continuous function. We say that $V : C \rightarrow \mathcal{R}$ is a Lyapunov function on a set $G \subset C$ relative to f if

V is continuous on $\bar{G}$, the closure of G , $\dot{V}$ is defined on G and $\dot{V} \leq 0$ on G .

Lemma 1(Hale [62]). Suppose $f(0) = 0$. Let V be a continuous functional defined on $C_H = C$ with $V(0) = 0$ and let $u(s)$ be a function, non-negative and continuous for $0 \leq s < \infty$, $u(s) \rightarrow \infty$ as $s \rightarrow \infty$ with $u(0) = 0$. If for all $\varphi \in C$, $u(|\phi(0)|) \leq V(\phi)$, $\dot{V}(\varphi) \leq 0$, then the zero solution of $\dot{x} = f(x_t)$ is stable.

If we define $Z = \{\varphi \in C_H : \dot{V}(\varphi) = 0\}$, then the zero solution of $\dot{x} = f(x_t)$ is asymptotically stable, provided that the largest invariant set in Z is $Q = 0$.

Besides, we consider the general non-autonomous delay differential system

$$\dot{x} = g(t, x_t), x_t = x(t + \theta), -r \leq \theta < 0, t \geq 0,$$

where $g : [0, \infty) \times C_\rho \rightarrow \mathcal{R}^n$ is a continuous mapping, $g(t, 0) = 0$, and we suppose that G takes closed bounded sets into bounded sets of $\mathcal{R}^n$. Here $(C, \|\cdot\|)$ is the Banach space of continuous function $\phi : [-r, 0] \rightarrow \mathcal{R}^n$ and for $\phi \in C$, $\|\phi\| = \sup_{-r \leq \theta \leq 0} |\phi(\theta)|$, $r > 0$, C_ρ is the open ball in C of radius ρ ; $C_\rho := \{\varphi \in C([-r, 0], \mathcal{R}^n) : \|\varphi\| < \rho\}$.

Theorem A(Yoshizawa [63 pp.191]). Assume that there is a Lyapunov functional $V_0(t, x)$ for $\dot{x} = g(t, x_t)$, and wedges satisfying;

(i) $W_1(|\phi(0)|) \leq V_0(t, \phi) \leq W_2(\|\phi\|)$, (where $W_1(r)$ and $W_2(r)$ are wedges),

(ii) $\dot{V}_0(t, \phi) \leq 0$.

Then the zero solution of $\dot{x} = g(t, x_t)$ is uniformly stable.

Let S be the set of $\phi \in C$ such that $\|\varphi\| > \rho$, denote by S^* the set of all functions $\phi \in C$ such that $|\phi(0)| \geq \rho$, where ρ may be large (Yoshizawa [63, pp.202]).

Theorem B (Yoshizawa [63, pp.202]). Suppose that there exists a continuous Lyapunov functional $V_0(t, \phi)$ defined for all $t \in \mathcal{R}^+$ and $\phi \in S^*$ which

satisfies the following conditions;

- (i) $a(|\phi(0)|) \leq V_0(t, \phi) \leq b_1(|\phi(0)|) + b_2(\|\phi\|)$,
 where $a(r), b_1(r), b_2(r) \in CI$, (CI denotes the families of continuous increasing functions), and are positive for $r > H$ and $a(r) - b_2(r) \rightarrow \infty$ as $r \rightarrow \infty$,
- (ii) $\dot{V}_0(t, \phi) \leq 0$. Then the solutions of $\dot{x} = g(t, x_t)$ are uniformly bounded.

Lemma 2. (Bellman [64, pp.288]). Let M be a real symmetric $n \times n$ -matrix. Then for any $X \in \mathcal{R}^n$

$$\delta_M \|X\|^2 \leq \langle MX, X \rangle \leq \Delta_M \|X\|^2,$$

where δ_M and Δ_M are, respectively, positive and simple, the least and greatest eigenvalues of the matrix M .

2. Stability

We introduce some basic assumptions needed in the proofs.

(A1) A is an $n \times n$ -symmetric constant matrix,

$$\delta_a \leq \lambda_i(A) \leq \Delta_a,$$

where δ_a and Δ_a are positive constants.

(A2) $H(0) = 0$, J_H exists and is an $n \times n$ -symmetric matrix,

$$H(X_1) \neq 0, \text{ when } X_1 \neq 0,$$

$$\delta_h \leq \lambda_i(J_H(X_1)) \leq \Delta_h \text{ for } X_1 \in \mathcal{R}^n,$$

where δ_h and Δ_h are positive constants.

(A3) $G(0) = 0$, J_G exists and is an $n \times n$ -symmetric matrix,

$$\delta_b \leq \lambda_i(J_G(X_2)) \leq \Delta_b \text{ for } X_2 \in \mathcal{R}^n,$$

where δ_b and Δ_b are positive constants, and the matrices J_G and J_H commute with each other.

Let $F(\cdot) \equiv 0$. The stability result of this paper is given by the following theorem.

Theorem 1. We assume that assumptions (A1) – (A3) hold. If

$$\tau_0 < \min \left\{ \frac{2(\delta_b - \beta \Delta_h)(1 - \tau_1)}{(\Delta_b + \Delta_h)(1 - \tau_1) + (\beta + 1)\Delta_b}, \frac{2(\beta \delta_b - 1)(1 - \tau_1)}{\beta(\Delta_b + \Delta_h)(1 - \tau_1) + (\beta + 1)\Delta_h} \right\},$$

then the zero solution of Eq. (1.2) is globally asymptotic stable, where τ_1 and β are positive constants with $0 < \tau_1 < 1$.

Proof. Define a functional $W_0 = W_0(t) = W_0(X_1(t), X_2(t), X_3(t))$ by

$$W_0 = 2 \int_0^1 \langle H(\sigma X_1), X_1 \rangle d\sigma + \langle AX_2, X_2 \rangle + 2\beta \int_0^1 \langle G(\sigma X_2), X_2 \rangle d\sigma$$

$$+ \beta \langle X_3, X_3 \rangle + 2 \langle X_2, X_3 \rangle + 2\beta \langle X_2, H(X_1) \rangle$$

$$(2.1) \quad + 2\lambda \int_{-\tau(t)}^0 \int_{t+s}^t \|X_2(\theta)\|^2 d\theta ds + 2\eta \int_{-\tau(t)}^0 \int_{t+s}^t \|X_3(\theta)\|^2 d\theta ds,$$

where β, λ and η are positive constants, the constants λ and η will be determined later in the proof.

From assumption (A₃), Lemma 2 and

$$G(0) = 0, \frac{\partial}{\partial \sigma} G(\sigma X_2) = J_G(\sigma X_2) X_2$$

We obtain

$$2 \int_0^1 \langle G(\sigma X_2), X_2 \rangle d\sigma = 2 \int_0^1 \int_0^1 \sigma_1 \langle J_G(\sigma_1 \sigma_2 X_2) X_2, X_2 \rangle d\sigma_1 d\sigma_2 \geq \delta_b \|X_2\|^2.$$

Similarly, it follows from

$$H(0) = 0, \frac{\partial}{\partial \sigma_1} H(\sigma_1 X_1) = J_H(\sigma_1 X_1) X_1$$

that

$$\frac{\partial}{\partial \sigma_1} \langle H(\sigma_1 X_1), H(\sigma_1 X_1) \rangle = 2 \langle J_H(\sigma_1 X_1) X_1, H(\sigma_1 X_1) \rangle.$$

Integrating, from $\sigma_1 = 0$ to $\sigma_1 = 1$, the both sides of the last estimates, respectively, we get

$$H(X_1) = \int_0^1 J_H(\sigma_1 X_1) X_1 d\sigma_1$$

and

$$\langle H(X_1), H(X_1) \rangle = 2 \int_0^1 \langle J_H(\sigma_1 X_1) X_1, H(\sigma_1 X_1) \rangle d\sigma_1.$$

It can also be seen that

$$2 \int_0^1 \langle H(\sigma_1 X_1), X_1 \rangle d\sigma_1 = 2 \int_0^1 \int_0^1 \sigma_1 \langle J_H(\sigma_1 \sigma_2 X_1) X_1, X_1 \rangle d\sigma_1 d\sigma_2.$$

Further, it is obvious that

$$\frac{\partial}{\partial \sigma_2} \langle H(\sigma_1 \sigma_2 X_1), J_H(\sigma_1 X_1) X_1 \rangle = \langle \sigma_1 J_H(\sigma_1 X_1) X_1, J_H(\sigma_1 X_1) X_1 \rangle.$$

Integrating the both sides of the last equality from $\sigma_2 = 0$ to $\sigma_2 = 1$, we obtain

$$\langle H(\sigma_1 X_1), J_H(\sigma_1 X_1) X_1 \rangle = \int_0^1 \langle \sigma_1 J_H(\sigma_1 X_1) X_1, J_H(\sigma_1 X_1) X_1 \rangle d\sigma_2.$$

Then, we have

$$\langle H(X_1), H(X_1) \rangle = 2 \int_0^1 \int_0^1 \langle \sigma_1 J_H(\sigma_1 X_1) X_1, J_H(\sigma_1 X_1) X_1 \rangle d\sigma_1 d\sigma_2.$$

From (2.1), the above discussion, Lemma 2 and the assumptions of Theorem 1, we can obtain

$$\begin{aligned}
 2W_0 &\geq 2 \int_0^1 \langle H(\sigma X_1), X_1 \rangle d\sigma + \langle AX_2, X_2 \rangle + \beta \delta_b \|X_2\|^2 \\
 &+ \beta \langle X_3, X_3 \rangle + 2 \langle X_2, X_3 \rangle + 2\beta \langle X_2, H(X_1) \rangle \\
 &= 2 \int_0^1 \langle H(\sigma X_1), X_1 \rangle d\sigma - \beta \delta_b^{-1} \langle H(X_1), H(X_1) \rangle \\
 &+ \beta \|\delta_b^{\frac{1}{2}} X_2 + \delta_b^{-\frac{1}{2}} H(X_1)\|^2 + \beta \|X_3 + \beta^{-1} X_2\|^2 \\
 &+ \langle (A - \beta^{-1} I) X_2, X_2 \rangle \\
 &\geq 2 \int_0^1 \langle H(\sigma X_1), X_1 \rangle d\sigma - \beta \delta_b^{-1} \langle H(X_1), H(X_1) \rangle \\
 &+ \beta \|X_3 + \beta^{-1} X_2\|^2 + \langle (A - \beta^{-1} I) X_2, X_2 \rangle \\
 &\geq 2 \int_0^1 \int_0^1 \sigma_1 \langle J_H(\sigma_1 \sigma_2 X_1) X_1, X_1 \rangle d\sigma_1 d\sigma_2 + \beta \|X_3 + \beta^{-1} X_2\|^2 \\
 &- 2\beta \delta_b^{-1} \int_0^1 \int_0^1 \langle \sigma_1 J_H(\sigma_1 X_1) X_1, J_H(\sigma_1 X_1) X_1 \rangle d\sigma_1 d\sigma_2 \\
 &+ \langle (A - \beta^{-1} I) X_2, X_2 \rangle \\
 &\geq (1 - \beta \delta_b^{-1} \Delta_h) \delta_h \|X_1\|^2 + \beta \|X_3 + \beta^{-1} X_2\|^2 + (\delta_a - \beta^{-1}) \|X_2\|^2.
 \end{aligned}$$

It is clear all the coefficients in the last inequality are positive and hence there exists a positive constant K such that

$$W_0 \geq K(\|X_1\|^2 + \|X_2\|^2 + \|X_3\|^2)$$

and

$$(2.2) \quad \|X_1\|^2 + \|X_2\|^2 + \|X_3\|^2 \leq K^{-1} W_0.$$

For the time derivative the functional W_0 by a straightforward calculation from (1.3) and (2.1), we obtain

$$\begin{aligned}
\dot{W}_0(t) = & -\langle (\beta A - I)X_3, X_3 \rangle - \langle G(X_2), X_2 \rangle + \beta \langle J_H(X_1)X_2, X_2 \rangle \\
& + \beta \int_{t-\tau(t)}^t \langle X_3(t), J_G(X_2(s))X_3(s) \rangle ds + \int_{t-\tau(t)}^t \langle X_2(s), J_G(X_2(s))X_3(s) \rangle ds \\
& + \beta \int_{t-\tau(t)}^t \langle X_3(t), J_H(X_1(s))X_2(s) \rangle ds + \int_{t-\tau(t)}^t \langle X_2(t), J_H(X_1(s))X_2(s) \rangle ds \\
& + \lambda \tau(t) \|X_2\|^2 + \eta \tau(t) \|X_3\|^2 - \lambda(1 - \tau'(t)) \int_{t-\tau(t)}^t \|X_2(\theta)\|^2 d\theta \\
& - \eta(1 - \tau'(t)) \int_{t-\tau(t)}^t \|X_3(\theta)\|^2 d\theta.
\end{aligned}$$

The assumptions of Theorem 1, $0 \leq \tau(t) \leq \tau_0$ and $\tau'(t) \leq \tau_1$, $0 < \tau_1 < 1$, imply that

$$\begin{aligned}
\langle (\beta A - I)X_3, X_3 \rangle & \geq (\beta \delta_a - 1) \|X_3\|^2, \\
\langle G(X_2), X_2 \rangle & = \int_0^1 \langle J_G(\sigma X_2)X_2, X_2 \rangle d\sigma \\
& \geq \int_0^1 \langle \delta_b X_2, X_2 \rangle d\sigma = \delta_b \|X_2\|^2, \\
\beta \langle J_H(X_1)X_2, X_2 \rangle & \leq \beta \Delta_h \|X_2\|^2,
\end{aligned}$$

$$\begin{aligned}
\beta \int_{t-\tau(t)}^t \langle X_3(t), J_G(X_2(s))X_3(s) \rangle ds & \leq \beta \|X_3(t)\| \int_{t-\tau(t)}^t \|J_G(X_2(s))\| \|X_3(s)\| ds \\
& \leq \beta \Delta_b \|X_3(t)\| \int_{t-\tau(t)}^t \|X_3(s)\| ds \\
& \leq \frac{1}{2} \beta \Delta_b \int_{t-\tau(t)}^t \left\{ \|X_3(t)\|^2 + \|X_3(s)\|^2 \right\} ds \\
& = \frac{1}{2} \beta \Delta_b \tau(t) \|X_3\|^2 + \frac{1}{2} \beta \Delta_b \int_{t-\tau(t)}^t \|X_3(s)\|^2 ds \\
& \leq \frac{1}{2} \beta \Delta_b \tau_0 \|X_3\|^2 + \frac{1}{2} \beta \Delta_b \int_{t-\tau(t)}^t \|X_3(s)\|^2 ds,
\end{aligned}$$

$$\begin{aligned}
 \int_{t-\tau(t)}^t \langle X_2(t), J_G(X_2(s))X_3(s) \rangle ds &\leq \|X_2(t)\| \int_{t-\tau(t)}^t \|J_G(X_2(s))\| \|X_3(s)\| ds \\
 &\leq \Delta_b \|X_2(t)\| \int_{t-\tau(t)}^t \|X_3(s)\| ds \\
 &\leq \frac{1}{2} \Delta_b \int_{t-\tau(t)}^t \left\{ \|X_2(t)\|^2 + \|X_3(s)\|^2 \right\} ds \\
 &= \frac{1}{2} \Delta_b \tau(t) \|X_2\|^2 + \frac{1}{2} \Delta_b \int_{t-\tau(t)}^t \|X_3(s)\|^2 ds \\
 &\leq \frac{1}{2} \Delta_b \tau_0 \|X_2\|^2 + \frac{1}{2} \Delta_b \int_{t-\tau(t)}^t \|X_3(s)\|^2 ds, \\
 \beta \int_{t-\tau(t)}^t \langle X_3(t), J_H(X_1(s))X_2(s) \rangle ds &\leq \beta \|X_3(t)\| \int_{t-\tau(t)}^t \|J_H(X_1(s))\| \|X_2(s)\| ds \\
 &\leq \beta \Delta_h \|X_3(t)\| \int_{t-\tau(t)}^t \|X_2(s)\| ds \\
 &\leq \frac{1}{2} \beta \Delta_h \int_{t-\tau(t)}^t \left\{ \|X_3(t)\|^2 + \|X_2(s)\|^2 \right\} ds \\
 &= \frac{1}{2} \beta \Delta_h \tau_0 \|X_3\|^2 + \frac{1}{2} \beta \Delta_h \int_{t-\tau(t)}^t \|X_2(s)\|^2 ds, \\
 \int_{t-\tau(t)}^t \langle X_2(t), J_H(X_1(s))X_2(s) \rangle ds &\leq \|X_2(t)\| \int_{t-\tau(t)}^t \|J_H(X_1(s))\| \|X_2(s)\| ds \\
 &\leq \Delta_h \|X_2(t)\| \int_{t-\tau(t)}^t \|X_2(s)\| ds \\
 &\leq \frac{1}{2} \Delta_h \int_{t-\tau(t)}^t \left\{ \|X_2(t)\|^2 + \|X_2(s)\|^2 \right\} ds \\
 &= \frac{1}{2} \Delta_h \tau(t) \|X_2\|^2 + \frac{1}{2} \Delta_h \int_{t-\tau(t)}^t \|X_2(s)\|^2 ds \\
 &\leq \frac{1}{2} \Delta_h \tau_0 \|X_2\|^2 + \frac{1}{2} \Delta_h \int_{t-\tau(t)}^t \|X_2(s)\|^2 ds, \\
 \lambda \tau(t) \|X_2\|^2 &\leq \lambda \tau_0 \|X_2\|^2, \\
 \eta \tau(t) \|X_3\|^2 &\leq \eta \tau_0 \|X_3\|^2, \\
 -\lambda(1 - \tau'(t)) \int_{t-\tau(t)}^t \|X_2(\theta)\|^2 d\theta &\leq -\lambda(1 - \tau_1) \int_{t-\tau(t)}^t \|X_2(\theta)\|^2 d\theta, \\
 -\eta(1 - \tau'(t)) \int_{t-\tau(t)}^t \|X_3(\theta)\|^2 d\theta &\leq -\eta(1 - \tau_1) \int_{t-\tau(t)}^t \|X_3(\theta)\|^2 d\theta.
 \end{aligned}$$

On gathering the obtained inequalities into $\dot{W}_0(t)$, we arrive at

$$\begin{aligned}\dot{W}_0(t) \leq & -\left\{\delta_b - \beta\Delta_h - \frac{1}{2}(\Delta_b + \Delta_h + 2\lambda)\tau_0\right\}\|X_2\|^2 \\ & -\left\{(\beta\delta_a - 1) - \frac{1}{2}(\beta(\Delta_b + \Delta_h) + 2\eta)\tau_0\right\}\|X_3\|^2 \\ & +\left\{\frac{1}{2}(\beta + 1)\Delta_h - \lambda(1 - \tau_1)\right\}\int_{t-\tau(t)}^t \|X_2(\theta)\|^2 ds \\ & +\left\{\frac{1}{2}(\beta + 1)\Delta_b - \eta(1 - \tau_1)\right\}\int_{t-\tau(t)}^t \|X_3(\theta)\|^2 ds.\end{aligned}$$

Let

$$\lambda = \frac{(\beta + 1)\Delta_h}{2(1 - \tau_1)}$$

and

$$\eta = \frac{(\beta + 1)\Delta_b}{2(1 - \tau_1)}.$$

Hence

$$\begin{aligned}\dot{W}_0(t) \leq & -\left\{\delta_b - \beta\Delta_h - \frac{1}{2}\left(\frac{(\Delta_b + \Delta_h)(1 - \tau_1) + (\beta + 1)\Delta_h}{1 - \tau_1}\right)\tau_0\right\}\|X_2\|^2 \\ & -\left\{(\beta\delta_b - 1) - \frac{1}{2}\left(\frac{\beta(\Delta_b + \Delta_h)(1 - \tau_1) + (\beta + 1)\Delta_b}{1 - \tau_1}\right)\tau_0\right\}\|X_3\|^2.\end{aligned}$$

If

$$\tau_0 < \min\left\{\frac{2(\delta_b - \beta\Delta_h)(1 - \tau_1)}{(\Delta_b + \Delta_h)(1 - \tau_1) + (\beta + 1)\Delta_b}, \frac{2(\beta\delta_b - 1)(1 - \tau_1)}{\beta(\Delta_b + \Delta_h)(1 - \tau_1) + (\beta + 1)\Delta_h}\right\},$$

$$0 < \tau_1 < 1,$$

then

$$\dot{W}_0(t) \leq -K_1\|X_2\|^2 - K_2\|X_3\|^2 \leq 0$$

for some positive constants K_1 and K_2 . In addition, we can easily see that

$$W_0(X_1, X_2, X_3) \rightarrow \infty \quad \text{as} \quad \|X_1\|^2 + \|X_2\|^2 + \|X_3\|^2 \rightarrow \infty.$$

Consider the set defined by

$$Q \equiv \{(X_1, X_2, X_3) : \dot{W}_0(X_1, X_2, X_3) = 0\}.$$

When we apply the LaSalle's invariance principle, we observe that $(X_1, X_2, X_3) \in Q$ implies that $X_2 = X_3 = 0$ and hence $X_1 = \mu$, ($\mu \neq 0$ is a constant vector). From the last estimate and system (1.3), we have $H(\mu) = 0$, which necessarily implies that $\mu = 0$ since $H(0) = 0$. Therefore

$$X_1 = X_2 = X_3 = 0.$$

In fact, this result implies that the largest invariant set contained in Q is $(0, 0, 0) \in Q$. By Lemma 1, we conclude that the zero solution of system (1.3) is asymptotically stable. Hence, the zero solution of Eq. (1.2) is the globally asymptotic stable. This completes the proof of Theorem 1.

Theorem 2. If assumptions (A1) – (A3) and

$$\tau_0 < \min \left\{ \frac{2(\delta_b - \beta\Delta_h)(1 - \tau_1)}{(\Delta_b + \Delta_h)(1 - \tau_1) + (\beta + 1)\Delta_b}, \frac{2(\beta\delta_b - 1)(1 - \tau_1)}{\beta(\Delta_b + \Delta_h)(1 - \tau_1) + (\beta + 1)\Delta_h} \right\},$$

hold, then the zero solution of Eq. (1.2) is uniformly stable, where τ_1 and β are positive constant with $0 < \tau_1 < 1$.

Proof. To prove Theorem 2, our main tool is the functional W_0 given by (2.1). It is clear from the proof of Theorem 1 that the functional W_0 and its time derivative satisfy the assumptions of Theorem A, except $W_2(\|\phi\|)$;

$$W_1(|\phi(0)|) \leq W_0 \leq W_2(\|\phi\|), \dot{W}(\phi) \leq 0.$$

Besides, subject to the assumptions of Theorem 2, it can be easily obtained that $W_0 \leq W_2(\|\phi\|)$. We omit the detail of the proof. This completes the proof of Theorem 2.

3. Boundedness

Let $F(\cdot) \neq 0$. The boundedness results of this paper are given by the following theorems.

Theorem 3. We assume that all the assumptions of Theorem 1 hold, except $F(\cdot) \equiv 0$. Further, we suppose that there exists a non-negative and continuous function $P = P(t)$ such that

$$\|F(t, X_1, X_2, X_3)\| \leq P(t) \text{ for all } t \geq 0, \max P(t) < \infty \text{ and } P \in L^1(0, \infty),$$

where $L^1(0, \infty)$ denotes the space of Lebesgue integrable functions.

If

$$\tau_0 < \min \left\{ \frac{2(\delta_b - \beta\Delta_h)(1 - \tau_1)}{(\Delta_b + \Delta_h)(1 - \tau_1) + (\beta + 1)\Delta_b}, \frac{2(\beta\delta_b - 1)(1 - \tau_1)}{\beta(\Delta_b + \Delta_h)(1 - \tau_1) + (\beta + 1)\Delta_h} \right\},$$

$$0 < \tau_1 < 1,$$

then there exists a constant $M > 0$ such that any solution $(X_1(t), X_2(t), X_3(t))$ of system (1.3) determined by

$$X_1(0) = X_{10}, X_2(0) = X_{20}, X_3(0) = X_{30}$$

satisfies

$$\|X_1(t)\| \leq M, \|X_2(t)\| \leq M, \|X_3(t)\| \leq M$$

for all $t \in \mathcal{R}^+$.

Proof. Let $F(\cdot) = F(t, X_1, X_2, X_3)$. For the case of $F(\cdot) \neq 0$, it can be concluded that

$$\dot{W}_0(t) \leq -K_1\|X_2\|^2 - K_2\|X_3\|^2 + \langle X_2, F(\cdot) \rangle + \langle \beta X_3, F(\cdot) \rangle.$$

Then

$$\dot{W}_0(t) \leq (\|X_2\| + \beta\|X_3\|)\|F(\cdot)\|$$

$$\leq K_3(\|X_2\| + \|X_3\|)\|F(\cdot)\|$$

$$\leq K_3(2 + \|X_2\|^2 + \|X_3\|^2)P(t)$$

$$\leq 2K_3P(t) + K^{-1}K_3W_0(t)P(t),$$

by the assumptions of Theorem 3 and (2.2), where

$$K_3 = \max\{1, \beta\}.$$

The integration of both sides of the last inequality, between 0 to t , ($t \geq 0$), leads that

$$W_0(t) \leq W_0(0) + 2K_3 \int_0^t P(s)ds + K^{-1}K_3 \int_0^t W_0(s)P(s)ds.$$

Let

$$M = W_0(0) + 2K_3 \int_0^\infty P(s)ds.$$

Then

$$W_0(t) \leq M + K^{-1}K_3 \int_0^\infty W_0(s)P(s)ds.$$

By noting the Gronwall-Bellman inequality (see Ahmad and Rama Mohana Rao [9, pp.41]), we can get

$$W_0(t) \leq M \exp(K^{-1}K_3 \int_0^\infty P(s)ds).$$

By the estimate $\|X_1\|^2 + \|X_2\|^2 + \|X_3\|^2 \leq K^{-1}W_0$ and the assumption $P \in L^1(0, \infty)$, we can conclude that all solutions of system (1.3) are bounded. This completes the proof of Theorem 3.

Theorem 4. If the assumptions Theorem 3 and

$$\tau_0 < \min \left\{ \frac{2(\delta_b - \beta\Delta_h)(1 - \tau_1)}{(\Delta_b + \Delta_h)(1 - \tau_1) + (\beta + 1)\Delta_b}, \frac{2(\beta\delta_b - 1)(1 - \tau_1)}{\beta(\Delta_b + \Delta_h)(1 - \tau_1) + (\beta + 1)\Delta_h} \right\},$$

hold, then the zero solution of Eq. (1.2) is uniformly bounded, where τ_1 is positive constant with $0 < \tau_1 < 1$.

Proof. To complete the poof of Theorem 4, the main tool is the functional W_0 given by (2.1). When we benefit from the functional W_0 and the assumptions of Theorem 4, we can easily complete the poof of Theorem 4. Therefore, we omit the details of the proof.

4. Conclusion

We consider a functional differential system of third order with variable delay. We investigate the globally asymptotic stability/uniformly stability/boundedness/ uniformly boundedness of solutions. The technique of proofs involves defining an appropriate Lyapunov functional. Our results include, improve and complete some recent results in the literature.

References

- [1] R. Reissig, G. Sansone, R. Conti, Non-linear differential equations of higher order. Translated from the German. Noordhoff International Publishing, Leyden, 1974.
- [2] A. T. Ademola, P. O. Arawomo, Stability and ultimate boundedness of solutions to certain third-order differential equations. *Appl. Math. E-Notes* 10, pp. 61-69, (2010).
- [3] A. T. Ademola, P. O. Arawomo, Stability and uniform ultimate boundedness of solutions of some third-order differential equations. *Acta Math. Acad. Paedagog. Nyhzi. (N.S.)* 27, No. 1, pp. 51-59, (2011).
- [4] A. T. Ademola, P. O. Arawomo, Asymptotic behaviour of solutions of third order nonlinear differential equations. *Acta Univ. Sapientiae Math.* 3, No. 2, pp. 197-211, (2011).
- [5] A. T. Ademola, P. O. Arawomo, Uniform stability and boundedness of solutions of nonlinear delay differential equations of the third order. *Math. J. Okayama Univ.* 55, pp. 157-166, (2013).
- [6] A. T. Ademola, B. S. Ogundare, M. O. Ogundiran, O. A. Adesina, Stability, boundedness, and existence of periodic solutions to certain third-order delay differential equations with multiple deviating arguments. *Int. J. Differ. Equ.*, Art. ID 213935, 12 pp., (2015).
- [7] A. U. Afuwape, J. E. Castellanos, Asymptotic and exponential stability of certain third- order non-linear delayed differential equations: frequency domain method. *Appl. Math. Comput.* 216, No. 3, pp. 940-950, (2010).
- [8] A. U. Afuwape, M. O. Omeike, Stability and boundedness of solutions of a kind of third- order delay differential equations. *Comput. Appl. Math.* 29, No. 3, pp. 329-342, (2010).
- [9] S. Ahmad, M. Rama Mohana Rao, Theory of ordinary differential equations. With applications in biology and engineering. Affiliated East-West Press Pvt. Ltd., New Delhi, (1999).

- [10] Y. Bai, C. Guo, New results on stability and boundedness of third order nonlinear delay differential equations. *Dynam. Systems Appl.* 22, No. 1, pp. 95-104, (2013).
- [11] J. O. C. Ezeilo, On the stability of solutions of certain differential equations of the third order. *Quart. J. Math. Oxford Ser. (2)* 11, pp. 64-69, (1960).
- [12] J. O. C. Ezeilo, H. O. Tjumla, Further results for a system of third order differential equations. *Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur. (8)* 58, No. 2, pp. 143-151, (1975).
- [13] J. R. Graef, D. Beldjerd, M. Remili, On stability, ultimate boundedness, and existence of periodic solutions of certain third order differential equations with delay. *PanAmer. Math. J.* 25, No. 1, pp. 82-94, (2015).
- [14] J. R. Graef, L. D. Oudjedi, M. Remili, Stability and square integrability of solutions of nonlinear third order differential equations. *Dyn. Contin. Discrete Impuls. Syst. Ser. A Math. Anal.* 22, No. 4, pp. 313-324, (2015).
- [15]] J. R. Graef, C. Tunc, Global asymptotic stability and boundedness of certain multidelay functional differential equations of third order. *Math. Methods Appl. Sci.* 38, No. 17, pp. 3747-3752, (2015).
- [16] E. Korkmaz, C. Tunc, Convergence to non-autonomous differential equations of second order. *J. Egyptian Math. Soc.* 23, No. 1, pp. 27-30, (2015).
- [17] A. M. Mahmoud, C. Tunc, Stability and boundedness of solutions of a certain n- dimensional nonlinear delay differential system of third-order. *Adv. Pure Appl. Math.* 7(1), pp. 1-11, (2016).
- [18] B. S. Ogundare, On boundedness and stability of solutions of certain third order delay differential equation. *J. Nigerian Math. Soc.* 31, pp. 55-68, (2012).
- [19] B. S. Ogundare, J. A. Ayanjinmi, O. A. Adesina, Bounded and L 2-solutions of certain third order non-linear differential equation with a square integrable forcing term. *Kragujevac J. Math.* 29, pp. 151-156, (2006).

- [20] A. L. Olutimo, Stability and ultimate boundedness of solutions of a certain third order nonlinear vector differential equation. *J. Nigerian Math. Soc.* 31, pp. 69-80, (2012).
- [21] M. O. Omeike, Stability and boundedness of solutions of a certain system of third order nonlinear delay differential equations. *Acta Univ. Palack. Olomuc. Fac. Rerum Natur. Math.* 54, No. 1, pp. 109-119, (2015).
- [22] C. Qian, On global stability of third-order nonlinear differential equations. *Nonlinear Anal.* 42, No. 4, Ser. A: Theory Methods, pp. 651-661, (2000).
- [23] M. Remili, L. D. Oudjedi, Stability and boundedness of the solutions of non-autonomous third order differential equations with delay. *Acta Univ. Palack. Olomuc. Fac. Rerum Natur. Math.* 53, No. 2, pp. 139-147, (2014).
- [24] A. I. Sadek, Stability and boundedness of a kind of third-order delay differential system. *Appl. Math. Lett.* 16, No. 5, 657-662, (2003).
- [25] K. E. Swick, Asymptotic behavior of the solutions of certain third order differential equations. *SIAM J. Appl. Math.* 19, pp. 96-102, (1970).
- [26] H. O. Tejumola, B. Tchegnani, Stability, boundedness and existence of periodic solutions of some third and fourth order nonlinear delay differential equations. *J. Nigerian Math. Soc.* 19, pp. 9-19, (2000).
- [27] C. Tunc, On the boundedness and periodicity of the solutions of a certain vector differential equation of third-order. Chinese translation in *Appl. Math. Mech.* 20, No. 2, pp. 153-160, (1999). *Appl. Math. Mech. (English Ed.)* 20, No. 2, pp. 163-170, (1999).
- [28] C. Tunc, Uniform ultimate boundedness of the solutions of third-order nonlinear differential equations. *Kuwait J. Sci. Engrg.* 32, No. 1, pp. 39-48, (2005).
- [29] C. Tunc, Boundedness of solutions of a third-order nonlinear differential equation. *JIPAM. J. Inequal. Pure Appl. Math.* 6, o. 1, Article 3, 6 pp., (2005).
- [30] C. Tunc, On the asymptotic behavior of solutions of certain third-order nonlinear differential equations. *J. Appl. Math. Stoch. Anal.*, No. 1, pp. 29-35, (2005).

- [31] C. Tunc, New results about stability and boundedness of solutions of certain non-linear third-order delay differential equations. Arab. J. Sci. Eng. Sect. A Sci. 31, No. 2, pp. 185-196, (2006).
- [32] C. Tunc, On the boundedness of solutions of certain nonlinear vector differential equations of third order. Bull. Math. Soc. Sci. Math. Roumanie (N. S.) 49 (97), No. 3, pp. 291-300, (2006).
- [33] C. Tunc, On the stability and boundedness of solutions to third order nonlinear differential equations with retarded argument. Nonlinear Dynam. 57, No. 1-2, pp. 97-106, (2009).
- [34] C. Tunc, On the stability and boundedness of solutions of nonlinear vector differential equations of third order. Nonlinear Anal. 70, No. 6, pp. 2232-2236, (2009).
- [35] C. Tunc, Bounded solutions to nonlinear delay differential equations of third order. Topol. Methods Nonlinear Anal. 34, No. 1, pp. 131-139, (2009).
- [36] C. Tunc, On the stability and boundedness of solutions of nonlinear third order differential equations with delay. Filomat 24, No. 3, pp. 1-10, (2010).
- [37] C. Tunc, Stability and bounded of solutions to non-autonomous delay differential equations of third order. Nonlinear Dynam. 62, No. 4, pp. 945-953, (2010).
- [38] C. Tunc, On some qualitative behaviors of solutions to a kind of third order nonlinear delay differential equations. Electron. J. Qual. Theory Differ. Equ., No. 12, 19 pp. , (2010).
- [39] C. Tunc, Stability and boundedness for a kind of non-autonomous differential equations with constant delay. Appl. Math. Inf. Sci. 7, No. 1, pp. 355-361, (2013).
- [40] C. Tunc, Stability and boundedness of the nonlinear differential equations of third order with multiple deviating arguments. Afr. Mat. 24, No. 3, pp. 381-390, (2013).
- [41] C. Tunc, On the qualitative behaviors of solutions of some differential equations of higher order with multiple deviating arguments. J. Franklin Inst. 351, No. 2, pp. 643-655, (2014).

- [42] C. Tunc, On the stability and boundedness of certain third order non-autonomous differential equations of retarded type. *Proyecciones* 34, No. 2, pp. 147-159, (2015).
- [43] C. Tunc, Global stability and boundedness of solutions to differential equations of third order with multiple delays. *Dynam. Systems Appl.* 24, pp. 467-478, (2015).
- [44] C. Tunc, Boundedness of solutions to certain system of differential equations with multiple delays. *Mathematical Modeling and Applications in Nonlinear Dynamics. Springer Book Series*, Chapter 5, pp. 109-123, (2016).
- [45] C. Tunc, M. Ate, Stability and boundedness results for solutions of certain third order nonlinear vector differential equations. *Nonlinear Dynam.* 45, No. 3-4, pp. 273-281, (2006).
- [46] C. Tunc, M. Gzen, Stability and uniform boundedness in multidelay functional differential equations of third order. *Abstr. Appl. Anal.*, Art. ID 248717, 7 pp. —, (2013).
- [47] C. Tunc, S. A. Mohammed, On the qualitative properties of differential equations of third order with retarded argument. *Proyecciones* 33, No. 3, pp. 325-347, (2014).
- [48] C. Tunc; E. Tunc, New ultimate boundedness and periodicity results for certain third- order nonlinear vector differential equations. *Math. J. Okayama Univ.* 48, pp. 159-172, (2006).
- [49] L. Zhang; L. Yu, Global asymptotic stability of certain third-order nonlinear differential equations. *Math. Methods Appl. Sci. Math. Methods Appl. Sci.* 36, No. 14, pp. 1845-1850, (2013).
- [50] Y. F. Zhu, On stability, boundedness and existence of periodic solution of a kind of third order nonlinear delay differential system. *Ann. Differential Equations* 8, No. 2, pp. 249-259, (1992).
- [51] A. U. Afuwape, P. Omari, F. Zanolin, Nonlinear perturbations of differential operators with nontrivial kernel and applications to third-order periodic boundary value problems. *J. Math. Anal. Appl.* 143, No. 1, pp. 35-56, (1989).

- [52] J. Andres, Periodic boundary value problem for certain nonlinear differential equations of the third order. *Math. Slovaca* 35, No. 3, 305-309, (1985).
- [53] K. O. Friderichs, On nonlinear vibrations of third order. *Studies in Nonlinear Vibration Theory*, pp. 65-103. Institute for Mathematics and Mechanics, New York University, (1946).
- [54] A. O. E. Animalu, J. O. C. Ezeilo, Some third order differential equations in physics. *Fundamental open problems in science at the end of the millennium*, Vol. III (Beijing, 1997), pp. 575-586, Hadronic Press, Palm Harbor, FL, (1999).
- [55] J. O. C. Ezeilo, J. Onyia, Nonresonant oscillations for some third-order differential equations. *J. Nigerian Math. Soc.* 3 (1984), 83-96 (1986).
- [56] K. E. Chlouverakis, J. C. Sprott, Chaotic hyperjerk systems. *Chaos Solitons Fractals* 28 (2006), no. 3, 739-746.
- [57] R. Eichhorn, S. J. Linz, P. Hnggi, Transformations of nonlinear dynamical systems to jerky motion and its application to minimal chaotic flows. *Phys Rev E* 58 (1998), 7151-7164.
- [58] S. J. Linz, On hyperjerky systems. *Chaos Solitons Fractals* 37 (2008), no. 3, 741-747.
- [59] J. Cronin-Scanlon, Some mathematics of biological oscillations. *SIAM Rev.* 19 (1977), no. 1, 100138.
- [60] L. L. Rauch, Oscillation of a third order nonlinear autonomous system. *Contributions to the Theory of Nonlinear Oscillations*, pp. 39-88. *Annals of Mathematics Studies*, no. 20. Princeton University Press, Princeton, N. J., (1950).
- [61] H. Smith, An introduction to delay differential equations with applications to the life sciences. *Texts in Applied Mathematics*, 57. Springer, New York, (2011).
- [62] J. Hale, Sufficient conditions for stability and instability of autonomous functional- differential equations. *J. Differential Equations* 1, pp. 452-482, (1965).

- [63] T. Yoshizawa, Stability theory by Liapunov's second method. Publications of the Mathematical Society of Japan, No. 9. The Mathematical Society of Japan, Tokyo (1966).
- [64] R. Bellman, Richard Introduction to matrix analysis. Reprint of the second edition 1970. With a foreword by Gene Golub. Classics in Applied Mathematics, 19. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, (1997).

Cemil Tunc

Department of Mathematics
Faculty of Science
Yuzuncu Yil University
65080, Van-Turkey
Turkey
e-mail : cemtunc@yahoo.com