

On I -statistically convergent sequence spaces defined by sequences of Orlicz functions using matrix transformation

Shyamal Debnath
Tripura University, India

and

Jayanta Debnath
National Institute of Technology, India
Received : November 2013. Accepted : May 2014

Abstract

Recently Savas and Das [12] introduced the notion of I -statistical convergence of sequences of real numbers. In this article we introduced the sequence spaces $W^{I(S)}(M, A, p)$, $W_0^{I(S)}(M, A, p)$ and $W_\infty^{I(S)}(M, A, p)$ of real numbers defined by I -statistical convergence using sequences of Orlicz function. We study some basic topological and algebraic properties of these spaces. We investigate some inclusion relations involving these spaces.

Key words : *Ideal, I -statistical convergence, Orlicz function, matrix transformation.*

AMS (2010) Classification No : *46A45, 40A35, 40C35.*

1. Introduction

The notion of statistical convergence was introduced by Fast [4] and Schoenberg [11], independently. Over the years and under different names statistical convergence has been discussed in the theory of Fourier analysis, ergodic theory and number theory by Buck [1], Esi and Et [3]. Moreover, statistical convergence is closely related to the concept of convergence in probability.

$$\text{i.e. } X_A(k) = \begin{cases} 1 & \text{if } k \in A \\ 0 & \text{if } k \in N \setminus A \end{cases} \quad \text{and} \quad d_n(A) = \frac{1}{n} \sum_{k=1}^n X_A(k)$$

The idea of statistical convergence depends on the density of subsets of the set N of natural numbers. Let N be the set of natural numbers. If $A \subseteq N$, then χ_A denotes the characteristic function of the set A

Then the number $\underline{d}(A) = \liminf d_n(A)$ and $\overline{d}(A) = \limsup d_n(A)$ are called the lower and upper asymptotic density of A respectively. If $\underline{d}(A) = \overline{d}(A) = d(A)$ then $d(A)$ is called the asymptotic density of A . We see that asymptotic density is limit of frequencies of numbers in the set $\{0, 1, 2, \dots\}$, therefore it is (when it exists) intuitively correct measure of size of subsets of integers. It is clear that any finite subset of N has natural density zero and $d(A^c) = 1 - d(A)$. Asymptotic density is (in some context) appropriate way to describe whether a subset of natural numbers is small or large.

A sequence $x = (x_n)$ is said to be statistically convergent to a number $L \in R$ if for each $\varepsilon > 0$, $d(A(\varepsilon)) = 0$, where $A(\varepsilon) = \{n \in N : |x_n - L| \geq \varepsilon\}$.

In recent years, generalizations of statistical convergence have appeared in the study of strong integral summability and the structure of ideals of bounded continuous functions on locally compact spaces. Kostyrko et.al.[7] presented a new generalization of statistical convergence and called it I -convergence. They used the notion of an ideal I of subsets of the set N to define such a concept.

Let X be a non-empty set. Then a family of sets $I \subset 2^X$ is said to be an ideal if I is additive, i.e, $A, B \in I \Rightarrow A \cup B \in I$ and hereditary i.e. $A \in I$, $B \subset A \Rightarrow B \in I$. A non-empty family of sets $F \subset 2^X$ is said to be a filter on X if and only if i) $\emptyset \notin F$ ii) for all $A, B \in F \Rightarrow A \cap B \in F$ iii) $A \in F$, $A \subset B \Rightarrow B \in F$. An ideal $I \subset 2^X$ is called non-trivial if $I \neq 2^X$. A non-trivial ideal I is called admissible iff $I \supset \{\{x\} : x \in X\}$. A non-trivial ideal I is maximal if there does not exist any non-trivial ideal $J \neq I$, containing I as a subset. For each ideal I there is a filter $F(I)$ corresponding to I i.e $F(I) = \{K \subseteq N : K^c \in I\}$, where $K^c = N - K$.

A sequence $x = (x_n)$ is said to be I -convergent to a number $L \in R$ if for a given $\varepsilon > 0$, we have $A(\varepsilon) = \{n \in N : |x_n - L| \geq \varepsilon\} \in I$. The element L is called the I -limit of the sequence $x = (x_n)$.

Example 1.1: Let $I = I_f = \{A \subseteq N : A \text{ is finite}\}$. Then I_f is nontrivial admissible ideal of N and the corresponding convergence coincides with ordinary convergence. If $I = I_d = \{A \subseteq N : d(A) = 0\}$, where $d(A)$ denotes the asymptotic density of the set A . Then I_d is a non-trivial admissible ideal of N and the corresponding convergence coincide with statistical convergence. For more on I -convergence one may refer to [2,16,19,20,21,23].

An Orlicz function $M : [0, \infty) \rightarrow [0, \infty)$ is a continuous, convex, non-decreasing function defined for $x > 0$ such that $M(0) = 0$ and $M(x) > 0$. If convexity of Orlicz function is replaced by $M(x + y) \leq M(x) + M(y)$, then this function is called modulus function. An Orlicz function M is said to satisfy Δ_2 -condition for all values of u , if there exists $K > 0$ such that $M(2u) \leq KM(u)$, $u \geq 0$. Let M be an Orlicz function which satisfies Δ_2 -condition and let $0 < \delta < 1$. Then for each $t \geq \delta$, we have $M(t) < K\delta^{-1}tM(2)$ for some constant $K > 0$. Two Orlicz functions M_1 and M_2 are said to be equivalent if there exists positive constants α, β and x_0 such that $M_1(\alpha) \leq M_2(x) \leq M_1(\beta)$, for all $0 \leq x < x_0$.

Lindenstrass and Tzafriri [8] studied some Orlicz type sequence spaces defined as follows:

$$\ell_M = \{(x_k) \in w : \sum M(\frac{|x_k|}{\rho}) < \infty, \text{ for } \rho > 0\}$$

The space ℓ_M with the norm

$$\|x\| = \inf\{\rho > 0 : \sum M(\frac{|x_k|}{\rho}) \leq 1\},$$

becomes a Banach space which is called an Orlicz sequence space. The space ℓ_M is closely related to ℓ_p which is an Orlicz sequence space with $M(t) = |t|^p$, for $1 \leq p < \infty$. Different classes of Orlicz sequence spaces were introduced and studied by Parasar and Choudhury [10], Esi and Et [3], Tripathy and Hazarika [17] and many others.

The notion of paranormed sequences was introduced by Nakano [9]. It was further investigated by Tripathy et. al. [14,15,22] and many others.

Definition 1.2: (Savas and Das[12]) A sequence $x = (x_k)$ is said to be I -statistically convergent to a number $L \in R$ if for each $\varepsilon > 0$, $\{n \in N : \frac{1}{n}|k \leq n : \|x_k - L\| \geq \varepsilon| \geq \delta\} \in I$.

The number L is called I -statistical limit of the sequence (x_k) and we write $I - st - \lim x_k = L$.

Remark 1.3: (Savas and Das[12]) Let $I = I_f = \{A \subseteq N : A \text{ is finite}\}$. Then I_f is nontrivial admissible ideal of N and I -statistical convergence coincides with statistical convergence.

Definition 1.4: A sequence space E is said to be solid (or normal) if $(y_k) \in E$ whenever $(x_k) \in E$ and $|y_k| \leq |x_k|$ for all $k \in N$.

Lemma 1.5: (One may refer to Kamthan and Gupta[6]) A sequence space E is normal implies that it is monotone.

Lemma 1.6: If $I \subset 2^N$ is a maximal ideal then for each $A \subset N$, we have either $A \in I$ or $N - A \in I$.

The following well-known inequality will be used throughout the article.

Let $p = (p_k)$ be any sequence of positive real numbers with $0 \leq p_k \leq \sup p_k = G$ and $D = \max\{1, 2^{G-1}\}$. Then $|a_k + b_k|^{p_k} \leq D(|a_k|^{p_k} + |b_k|^{p_k})$ for all $k \in N$ and $a_k, b_k \in C$.

Also $|a_k|^{p_k} \leq \max\{1, |a|^G\}$ for all $a \in C$.

2. Main Result

Let $M = (M_k)$ be a sequence of Orlicz functions and $A = (a_{ik})$ be an infinite matrix and $x = (x_k)$ be a sequence of real or complex numbers. We write $Ax = (A_k(x))$ if $A_k(x) = \sum_k a_{ik} x_k$ converges for each i .

We define the following sequence spaces in this article:

$W^{I(S)}(M, A, p) = \{(x_k) \in w : \{n \in N : \frac{1}{n} |\{k \leq n : \sum_{k=1}^n [M_k(\frac{\|A_k(x) - L\|}{\rho})]^{p_k} \geq \varepsilon\}| \geq \delta\} \in I \text{ for some } \rho > 0 \text{ and } L \in R\}$.

$W_0^{I(S)}(M, A, p) = \{(x_k) \in w : \{n \in N : \frac{1}{n} |\{k \leq n : \sum_{k=1}^n [M_k(\frac{\|A_k(x)\|}{\rho})]^{p_k} \geq \varepsilon\}| \geq \delta\} \in I \text{ for some } \rho > 0\}$.

$W_\infty^{I(S)}(M, A, p) = \{(x_k) \in w : \{n \in N : \frac{1}{n} |\{k \leq n : \sum_{k=1}^n [M_k(\frac{\|A_k(x)\|}{\rho})]^{p_k} \geq M\}| \geq \delta\} \in I \text{ for some } M > 0\}$.

$W_\infty(M, A, p) = \{(x_k) \in w : \{n \in N : \sup \frac{1}{n} \sum_{k=1}^n [M_k(\frac{\|A_k(x)\|}{\rho})]^{p_k} < \infty\}\}$.

From the above definition it is obvious that

$W_0^{I(S)}(M, A, p) \subset W^{I(S)}(M, A, p) \subset W_\infty^{I(S)}(M, A, p)$.

Theorem 2.1: The spaces $W_0^{I(S)}(M, A, p)$, $W^{I(S)}(M, A, p)$ and $W_\infty^{I(S)}(M, A, p)$ are linear space.

Proof: We prove the result for the space $W_0^{I(S)}(M, A, p)$. The other result can be established in similar way.

Let $x = (x_k)$ and $y = (y_k)$ be any two elements in $W_0^{I(S)}(M, A, p)$. Then there exists $\rho_1 > 0$ and $\rho_2 > 0$ such that

$$A = \{n \in N : \frac{1}{n} |\{k \leq n : \sum_{k=1}^n [M_k(\frac{\|A_k(x)\|}{\rho_1})]^{p_k} \geq \frac{\varepsilon}{2}\}| \geq \delta\} \in I$$

$$\text{and } B = \{n \in N : \frac{1}{n} |\{k \leq n : \sum_{k=1}^n [M_k(\frac{\|A_k(x)\|}{\rho_2})]^{p_k} \geq \frac{\varepsilon}{2}\}| \geq \delta\} \in I$$

Let a, b be any scalars. By the continuity of the sequence $M = (M_k)$ the following inequality holds:

$$\begin{aligned} \sum_{k=1}^n [M_k(\frac{\|A_k(ax+by)\|}{|a|\rho_1+|b|\rho_2})]^{p_k} &\leq DK \sum_{k=1}^n [M_k(\frac{\|A_k(x)\|}{\rho_1})]^{p_k} + DK \sum_{k=1}^n [M_k(\frac{\|A_k(y)\|}{\rho_2})]^{p_k} \\ &\leq D \sum_{k=1}^n [\frac{|a|}{|a|\rho_1+|b|\rho_2} M_k(\frac{\|A_k(x)\|}{\rho_1})]^{p_k} + D \sum_{k=1}^n [\frac{|b|}{|a|\rho_1+|b|\rho_2} M_k(\frac{\|A_k(x)\|}{\rho_1})]^{p_k} \end{aligned}$$

where $K = \max\{1, \frac{|a|}{|a|\rho_1+|b|\rho_2}, \frac{|b|}{|a|\rho_1+|b|\rho_2}\}$.

From the above relation we get the following:

$$\{n \in N : \frac{1}{n} |\{k \leq n : \sum_{k=1}^n [M_k(\frac{\|A_k(ax+by)\|}{|a|\rho_1+|b|\rho_2})]^{p_k} \geq \frac{\varepsilon}{2}\}| \geq \delta\} \subseteq$$

$$\{n \in N : \frac{1}{n} |\{k \leq n : \sum_{k=1}^n DK [M_k(\frac{\|A_k(x)\|}{\rho_1})]^{p_k} \geq \frac{\varepsilon}{2}\}| \geq \delta\}$$

$$\cup \{n \in N : \frac{1}{n} |\{k \leq n : \sum_{k=1}^n DK [M_k(\frac{\|A_k(y)\|}{\rho_2})]^{p_k} \geq \frac{\varepsilon}{2}\}| \geq \delta\}$$

This completes the proof.

Theorem 2.2: The space $W_\infty(M, A, p)$ is a paranormed spaces (not totally paranormed) with the paranorm g defined by:

$$g(x) = \inf\{\rho^{\frac{p_k}{H}} : \sup_k M_k(\frac{\|A_k(x)\|}{\rho}) \leq 1, \text{ for } \rho > 0\}, \text{ where } H = \max\{1, \sup_k p_k\}.$$

Proof: It is obvious that $g(\theta) = 0$ (where θ is the sequence of zeros), $g(-x) = g(x)$ and it can be easily shown that $g(x+y) \leq g(x) + g(y)$.

Let $t_n \rightarrow L$, where $t_n, L \in C$ and let $g(x_n - x) \rightarrow 0$, as $n \rightarrow \infty$. To prove that $g(t_n x_n - Lx) \rightarrow 0$, as $n \rightarrow \infty$. We put

$$A = \{\rho_1 > 0 : \sup_k [M_k(\frac{\|A_k(x)\|}{\rho_1})]^{p_k} \leq 1\}$$

and

$$B = \{\rho_2 > 0 : \sup_k [M_k(\frac{\|A_k(x)\|}{\rho_2})]^{p_k} \leq 1\}$$

By the continuity of the sequence $M = (M_k)$, we observe that

$$\begin{aligned} M_k(\frac{\|A_k(t_n x_n - Lx)\|}{|t_n - L|\rho_1 + |L|\rho_2}) &\leq M_k(\frac{\|A_k(t_n x_n - Lx_n)\|}{|t_n - L|\rho_1 + |L|\rho_2}) + M_k(\frac{\|A_k(Lx_n - Lx)\|}{|t_n - L|\rho_1 + |L|\rho_2}) \\ &\leq \frac{|t_n - L|\rho_1}{|t_n - L|\rho_1 + |L|\rho_2} M_k(\frac{\|A_k(x_n)\|}{\rho_1}) + \frac{|L|\rho_2}{|t_n - L|\rho_1 + |L|\rho_2} M_k(\frac{\|A_k(x_n - x)\|}{\rho_2}) \end{aligned}$$

From the above inequality it follows that

$$\sup_k [M_k(\frac{\|A_k(t_n x_n - Lx)\|}{|t_n - L|\rho_1 + |L|\rho_2})]^{p_k} \leq 1$$

and hence

$$g(t_n x_n - Lx) = \inf\{(|t_n - L|\rho_1 + |L|\rho_2)^{\frac{p_k}{H}} : \rho_1 \in A, \rho_2 \in B\}$$

$$\leq (|t_n - L|)^{\frac{p_k}{H}} \inf\{\rho_1^{\frac{p_k}{H}} : \rho_1 \in A\} + (|L|)^{\frac{p_k}{H}} \inf\{\rho_2^{\frac{p_k}{H}} : \rho_2 \in B\}$$

$$\leq \max\{|t_n - L|, (|t_n - L|)^{\frac{p_k}{H}}\} g(x_n) + \max\{|L|, (|L|)^{\frac{p_k}{H}}\} g(x_n - x)$$

As $g(x_n) \leq g(x) + g(x_n - x)$ for all $n \in N$, hence the right hand side of the above relation tends to zero as $n \rightarrow \infty$.

This completes the proof.

Proposition 2.3: Let $M = (M_k)$ and $N = (N_k)$ be sequences of Orlicz functions. Then the following hold:

(i) $W_0^{I(S)}(N, A, p) \subseteq W_0^{I(S)}(MoN, A, p)$, provided $p = (p_k)$ such that $G_0 = \inf p_k > 0$.

(ii) $W_0^{I(S)}(M, A, p) \cap W_0^{I(S)}(N, A, p) \subseteq W_0^{I(S)}(M + N, A, p)$.

Theorem 2.4: The spaces $W_0^{I(S)}(M, A, p)$ and $W^{I(S)}(M, A, p)$ are normal and monotone.

Proof: Let $x = (x_k) \in W_0^{I(S)}(M, A, p)$ and $y = (y_k)$ be such that $|y_k| \leq |x_k|$. Then for $\varepsilon > 0$,

$$\{n \in N : \frac{1}{n} |\{k \leq n : \sum_{k=1}^n [M_k(\frac{\|A_k(x)\|}{\rho})]^{p_k} \geq \varepsilon\}| \geq \delta\}$$

$$\supseteq \{n \in N : \frac{1}{n} |\{k \leq n : \sum_{k=1}^n [M_k(\frac{\|A_k(y)\|}{\rho})]^{p_k} \geq \varepsilon\}| \geq \delta\} \in I.$$

The result follows from the above relation. Thus the space $W_0^{I(S)}(M, A, p)$ is normal and hence monotone by lemma 1.5. Similarly for the other.

Proposition 2.5: Let $0 < p_k \leq q_k$ and $\frac{q_k}{p_k}$ be bounded. Then $W_0^{I(S)}(M, A, q) \subseteq W_0^{I(S)}(M, A, p)$

Proposition 2.6: For any two sequence $p = (p_k)$ and $q = (q_k)$ of positive real numbers, the following hold:

$$Z(M, A, p) \cap Z(M, A, q) \neq \emptyset \text{ for } Z = W^{I(S)}, W_0^{I(S)}, W_\infty^{I(S)}.$$

References

- [1] Buck R. C., The measure theoretic approach to density, Amer. J. Math., 68, pp. 560-580, (1946).
- [2] Debnath S. and Debnath J., Some generalized statistical convergent sequence spaces of fuzzy numbers via ideals, Math. Sci. Lett., 2, No. 2, pp. 151-154, (2013).
- [3] Esi A. and Et M., Some new spaces defined by Orlicz functions, Indian J. Pure and Appl. Math., 31 (8), pp. 967-972, (2000).
- [4] Fast H., Sur la convergence statistique, Colloq. Math., pp. 2241-244, (1951).

- [5] Fridy J. A., On statistical convergence, *Analysis*, pp. 301-313, (1985).
- [6] Kamthan P. K. and Gupta M., *Sequence spaces and series* (1980).
- [7] Kostyrko P. Salat T., and Wilczynski W., I-convergence, *Real analysis exchange*, 26 (2), pp. 669-686, (2000/2001).
- [8] Lindenstrauss J. and Tzafriri L., On Orlicz sequence spaces, *Israel J. Math.*, 101, pp. 379-390, (1971).
- [9] Nakano H., Modular sequence spaces, *Proc. Japan Acad.*, 27, pp. 508-512, (1951).
- [10] Parashar S.D and Choudhury B., Sequence space defined by orlicz functions, *Indian J. Pure and Appl. Math.*, 25(14), pp. 419-428, (1994).
- [11] Salat T., On statistically convergent sequences of real numbers, *Math. Slovaca*, 30, pp. 139-150, (1980).
- [12] Savas E. and Das P., A generalized statistical convergence via ideals, *Applied mathematics letters*, 24, pp. 826-830, (2011).
- [13] Schoenberg I. J., The integrability of certain functions and related summability methods, *Am. Math. Mon.*, 66, pp. 361-375, (1951).
- [14] Tripathy B. C. and Chandra P., On some generalized difference paranormed sequence spaces associated with multiplier sequences defined by modulus function, *Anal. Theory Appl.*, 27 (1), pp. 21-27, (2011).
- [15] Tripathy B. C. and Dutta H., On some new paranormed difference sequence spaces defined by Orlicz functions, *Kyungpook Mathematical Journal*, 50 (1), pp. 59-69, (2010).
- [16] Tripathy B. C., and Dutta A. J., On I-acceleration convergence of sequences of fuzzy real numbers, *Math. Modell. Analysis*, 17 (4), pp. 549-557, (2012).
- [17] Tripathy B. C. and Hazarika B., Some I-convergent sequence spaces defined by orlicz functions, *Acta Math. Appl. Sin.*, 27(1), pp. 149-154, (2011).
- [18] Tripathy B. C. and Hazarika B., Paranorm I-convergent sequence spaces, *Math. Slovaca*, 59 (4), pp. 485-494, (2009).

- [19] Tripathy B. C. and Hazarika B., I-convergent sequence spaces associated with multiplier sequence spaces, *Mathematical Inequalities and Applications*, 11 (3), pp. 543-548, (2008).
- [20] Tripathy B. C. and Hazarika B., I-monotonic and I-convergent sequences, *Kyungpook Math. Journal*, 51 (2), pp. 233-239, (2011).
- [21] Tripathy B. C. and Mahanta S., On I-acceleration convergence of sequences, *Journal of the Franklin Institute*, 347, pp. 591-598, (2010).
- [22] Tripathy B. C. and Sen M., Characterization of some matrix classes involving paranormed sequence spaces, *Tamkang Jour. Math.*, 37(2), pp. 155-162, (2006).
- [23] Tripathy B.C., Sen M., and Nath S., I-convergence in probabilistic n -normed space, *Soft Comput.*, 16, pp. 1021-1027, (2012), DOI 10.1007/s00500-011-0799-8.

Shyamal Debnath

Department of Mathematics,
Tripura University,
Agartala-799022,
India
e-mail : shyamalnitamath@gmail.com

and

Jayanta Debnath

Department of Mathematics,
National Institute of Technology,
Agartala-799055,
India
e-mail : mailme_jdebnath@rediffmail.com