

## Some Umbral Calculus Presentations of the Chan-Chyan-Srivastava Polynomials and the Erkuş-Srivastava Polynomials

*H. M. Srivastava*

*University of Victoria, Canada*

*K. S. Nisar*

*Salman Bin Abdu-Aziz University, Saudi Arabia  
and*

*Mumtaz Ahmad Khan*

*Aligarh Muslim University, India*

*Received : February 2013. Accepted : March 2013*

### Abstract

*In their recent investigation involving differential operators for the generalized Lagrange polynomials, Chan et. al. [3] encountered and proved a certain summation identity and several other results for the Lagrange polynomials in several variables, which are popularly known in the literature as the Chan-Chyan-Srivastava polynomials. These multivariable polynomials have been studied systematically and extensively in the literature ever since then (see, for example, [1], [4], [9], [11], [12] and [13]). In the present paper, we investigate umbral calculus presentations of the Chan-Chyan-Srivastava polynomials and also of their substantially more general form, the Erkuş-Srivastava polynomials [9]. Some other closely-related results are also considered.*

**2010 Mathematics Subject Classification :** *Primary 33C45, 33C65; Secondary 33C05.*

**Key Words and Phrases :** *Lagrange polynomials; Hermite-Kampé de Fériet polynomials; Lagrange-Hermite polynomials; Chan-Chyan-Srivastava polynomials; Erkuş-Srivastava polynomials; Umbral calculus; Pochhammer symbol; Multinomial theorem and multinomial coefficients; Principle of monoumbrality; Monoumbral expansions.*

## 1. Introduction, Definitions and Notations

The familiar (two-variable) polynomials  $g_n^{(\alpha, \beta)}(x, y)$ , which are generated by

$$(1 - xz)^{-\alpha} (1 - yz)^{-\beta} = \sum_{n=0}^{\infty} g_n^{(\alpha, \beta)}(x, y) z^n \quad (1.1)$$

$$(\alpha, \beta \in \mathbf{C}; |z| < \min\{|x|^{-1}, |y|^{-1}\}),$$

are known as the Lagrange polynomials which occur in certain problems in statistics (*cf.*, *e.g.*, Erdélyi *et al.* [8, p. 267]; see also [15, p. 441 *et seq.*]).

The (three-variable) Lagrange polynomials  $g_n^{(\alpha, \beta, \gamma)}(x, y, z)$ , which are defined by means of the generating function:

$$(1 - xt)^{-\alpha} (1 - yt)^{-\beta} (1 - zt)^{-\gamma} = \sum_{n=0}^{\infty} g_n^{(\alpha, \beta, \gamma)}(x, y, z) \frac{t^n}{n!} \quad (1.2)$$

$$(\alpha, \beta, \gamma \in \mathbf{C}; |t| < \min\{|x|^{-1}, |y|^{-1}, |z|^{-1}\}),$$

were studied recently by Khan and Shukla [10]. Subsequently, Chan et al. [3] introduced and investigated the multivariable extension of the classical Lagrange polynomials  $g_n^{(\alpha, \beta)}(x, y)$  generated by (1.1).

These multivariable Lagrange polynomials  $g_n^{(\alpha_1, \dots, \alpha_r)}(x_1, \dots, x_r)$ , which are popularly known in the literature as the *Chan-Chyan-Srivastava polynomials*, are generated by (see, for details, [3]; see also [4], [12] and [13])

$$\prod_{j=1}^r \{(1 - x_j z)^{-\alpha_j}\} = \sum_{n=0}^{\infty} g_n^{(\alpha_1, \dots, \alpha_r)}(x_1, \dots, x_r) z^n \quad (1.3)$$

$$(\alpha_j \in \mathbf{C} \quad (j = 1, \dots, r); |z| < \min\{|x_1|^{-1}, \dots, |x_r|^{-1}\}),$$

so that, upon comparison with the generating function (1.2), we have the following relationship:

$$g_n^{(\alpha, \beta, \gamma)}(x, y, z) = n! g_n^{(\alpha, \beta, \gamma)}(x, y, z)$$

$$(n \in \mathbf{N}_0 := \mathbf{N} \cup \{0\}; \mathbf{N} := \{1, 2, 3, \dots\})$$

with the (three-variable) Lagrange polynomials  $g_n^{(\alpha, \beta, \gamma)}(x, y, z)$  studied by Khan and Shukla [10].

Clearly, the defining generating function (1.3) yields the explicit representation given by [3, p. 140, Eq. (6)]

$$\begin{aligned}
 g_n^{(\alpha_1, \dots, \alpha_r)}(x_1, \dots, x_r) &= \sum_{\substack{k_1, \dots, k_r \in \mathbf{N}_0 \\ (k_1 + \dots + k_r = n)}} (\alpha_1)_{k_1} \cdots (\alpha_r)_{k_r} \frac{x_1^{k_1}}{k_1!} \cdots \frac{x_r^{k_r}}{k_r!} \\
 &= \sum_{k_1, \dots, k_{r-1}=0}^{k_1 + \dots + k_{r-1} \leq n} \frac{(\alpha_1)_{n-k_1-\dots-k_{r-1}} (\alpha_2)_{k_1} \cdots (\alpha_r)_{k_{r-1}}}{(n-k_1-\dots-k_{r-1})! k_1! \cdots k_{r-1}!} \\
 &\quad \cdot x_1^{n-k_1-\dots-k_{r-1}} x_2^{k_1} \cdots x_r^{k_{r-1}}
 \end{aligned}
 \tag{1.4}$$

or, equivalently, by [11, p. 522, Eq. (17)]

$$\begin{aligned}
 g_n^{(\alpha_1, \dots, \alpha_r)}(x_1, \dots, x_r) &= \sum_{n_{r-1}=0}^n \sum_{n_{r-2}=0}^{n_{r-1}} \cdots \sum_{n_1=0}^{n_2} \frac{(\alpha_1)_{n_1} (\alpha_2)_{n_2-n_1} \cdots (\alpha_r)_{n-n_{r-1}}}{n_1! (n_2-n_1)! \cdots (n-n_{r-1})!} \\
 &\quad \cdot x_1^{n_1} x_2^{n_2-n_1} \cdots x_r^{n-n_{r-1}} \\
 &= \sum_{k_1=0}^n \sum_{k_2=0}^{k_1} \cdots \sum_{k_{r-1}=0}^{k_{r-2}} \frac{(\alpha_1)_{n-k_1} (\alpha_2)_{k_2-k_1} \cdots (\alpha_r)_{k_{r-1}}}{(n-k_1)! (k_2-k_1)! \cdots k_{r-1}!} \\
 &\quad \cdot x_1^{n-k_1} x_2^{k_2-k_1} \cdots x_r^{k_{r-1}},
 \end{aligned}
 \tag{1.5}$$

where, as usual,  $(\lambda)_n$  denotes the Pochhammer symbol given by

$$(\lambda)_0 := 1 \quad \text{and} \quad (\lambda)_n := \lambda(\lambda+1) \cdots (\lambda+n-1) \quad (n \in \mathbf{N}).$$

Altın and Erkuş [1] presented a multivariable extension of the so-called Lagrange-Hermite polynomials generated by (see [1, p. 239, Eq. (1.2)])

$$\begin{aligned}
 (1.6) \quad \prod_{j=1}^r \{(1 - x_j z^j)^{-\alpha_j}\} &= \sum_{n=0}^{\infty} h_n^{(\alpha_1, \dots, \alpha_r)}(x_1, \dots, x_r) z^n \\
 &\quad \left( \alpha_j \in \mathbf{C} \quad (j = 1, \dots, r); |z| < \min_{j \in \{1, \dots, r\}} \{|x_j|^{-1/j}\} \right).
 \end{aligned}$$

The case  $r = 2$  of the polynomials given by (1.6) corresponds to the familiar (two-variable) Lagrange-Hermite polynomials considered by Dattoli *et al.* [6].

The multivariable (Erkuş-Srivastava) polynomials

$\mathcal{U}_{n;\ell_1,\dots,\ell_r}^{(\alpha_1,\dots,\alpha_r)}(x_1,\dots,x_r)$ , which are defined by the following generating function [9, p. 268, Eq. (3)]:

$$(1.7) \quad \prod_{j=1}^r \{(1 - x_j z^{\ell_j})^{-\alpha_j}\} = \sum_{n=0}^{\infty} \mathcal{U}_{n;\ell_1,\dots,\ell_r}^{(\alpha_1,\dots,\alpha_r)}(x_1,\dots,x_r) z^n$$

$$(\alpha_j \in \mathbf{C} \quad (j = 1, \dots, r); \ell_j \in \mathbf{N} \quad (j = 1, \dots, r);$$

$$|z| < \min\{|x_1|^{-1/\ell_1}, \dots, |x_r|^{-1/\ell_r}\},$$

are a unification (*and* generalization) of several known families of multivariable polynomials including (for example) the Chan-Chyan-Srivastava polynomials  $g_n^{(\alpha_1,\dots,\alpha_r)}(x_1,\dots,x_r)$  defined by (1.3) (see, for details, [9]). Obviously, the Chan-Chyan-Srivastava polynomials  $g_n^{(\alpha_1,\dots,\alpha_r)}(x_1,\dots,x_r)$  follow as the special case of the Erkuş-Srivastava polynomials  $\mathcal{U}_{n;\ell_1,\dots,\ell_r}^{(\alpha_1,\dots,\alpha_r)}(x_1,\dots,x_r)$  when  $\ell_j = 1$  ( $j = 1, \dots, r$ ).

Moreover, the Lagrange-Hermite polynomials  $h_n^{(\alpha_1,\dots,\alpha_r)}(x_1,\dots,x_r)$  follow as a special case of the Erkuş-Srivastava polynomials  $\mathcal{U}_{n;\ell_1,\dots,\ell_r}^{(\alpha_1,\dots,\alpha_r)}(x_1,\dots,x_r)$  when  $\ell_j = j$  ( $j = 1, \dots, r$ ).

The generating function (1.7) yields the following *explicit* representation [9, p. 268, Eq. (4)]:

$$(1.8) \quad \mathcal{U}_{n;\ell_1,\dots,\ell_r}^{(\alpha_1,\dots,\alpha_r)}(x_1,\dots,x_r) = \sum_{\substack{k_1,\dots,k_r \in \mathbf{N}_0 \\ (\ell_1 k_1 + \dots + \ell_r k_r = n}} (\alpha_1)_{k_1} \cdots (\alpha_r)_{k_r} \frac{x_1^{k_1}}{k_1!} \cdots \frac{x_r^{k_r}}{k_r!},$$

which, in the special case when  $\ell_j = 1$  ( $j = 1, \dots, r$ ), corresponds to the *first* expression in (1.4).

Each of the above families of multivariable polynomials has been investigated systematically and extensively in the literature ever since the publication of the work by Chan *et al.* [9] (see, for example, [1], [4], [9], [11], [12] and [13]). The main objective of the present paper is to derive umbral calculus presentations of the Chan-Chyan-Srivastava polynomials  $g_n^{(\alpha_1,\dots,\alpha_r)}(x_1,\dots,x_r)$  generated by (1.3) and also of the substantially more general Erkuş-Srivastava polynomials  $\mathcal{U}_{n;\ell_1,\dots,\ell_r}^{(\alpha_1,\dots,\alpha_r)}(x_1,\dots,x_r)$  generated by (1.7). Upon suitable specialization, this last umbral calculus presentation is shown to yield the corresponding result for the polynomials  $h_n^{(\alpha_1,\dots,\alpha_r)}(x_1,\dots,x_r)$  generated by (1.6).

## 2. Umbral Calculus Presentations

The Chan-Chyan-Srivastava polynomials in (1.3) exhibits a structure which, according to the prescription provided in [7], can be viewed as the umbral image of ordinary monomials. Indeed, by using the generating function in (1.3) and an elementary integral identity in the form [2] :

$$(2.1) \quad \kappa^{-\nu} = \frac{1}{\Gamma(\nu)} \int_0^\infty e^{-\kappa t} t^{\nu-1} dt \quad \left( \min\{\Re(\kappa), \Re(\nu)\} > 0 \right),$$

we can easily derive the following integral representation of the Chan-Chyan-Srivastava polynomials:

$$(2.2) \quad g_n^{(\alpha_1, \dots, \alpha_r)}(x_1, \dots, x_r) = \frac{1}{\Gamma(\alpha_1) \cdots \Gamma(\alpha_r)} \int_0^\infty \cdots \int_0^\infty e^{-(\xi_1 + \cdots + \xi_r)} \cdot \xi_1^{\alpha_1-1} \cdots \xi_r^{\alpha_r-1} \mathcal{P}_n(x_1 \xi_1, \dots, x_r \xi_r) d\xi_1 \cdots d\xi_r,$$

where

$$(2.3) \quad \mathcal{P}_n(x_1, \dots, x_r) := \frac{(x_1 + \cdots + x_r)^n}{n!} = \frac{1}{n!} \sum_{\substack{k_1, \dots, k_r \in \mathbf{N}_0 \\ (k_1 + \cdots + k_r = n)}} \binom{n}{k_1, \dots, k_r} x_1^{k_1} \cdots x_r^{k_r}$$

in terms of the multinomial coefficients given by

$$\binom{n}{k_1, \dots, k_r} := \frac{n!}{k_1! \cdots k_r!} \quad (n, k_1, \dots, k_r \in \mathbf{N}_0).$$

Alternatively, the multinomial theorem (see, for example, [15, p. 87, Problem 5]) used in (2.3) can indeed be restated as follows:

$$(2.4) \quad \mathcal{P}_n(x_1, \dots, x_r) = \frac{1}{n!} \sum_{k_1=0}^n \sum_{k_2=0}^{n-k_1} \cdots \sum_{k_{r-1}=0}^{n-k_1-\cdots-k_{r-2}} \binom{n}{k_1} \binom{n-k_1}{k_2} \cdots \binom{n-k_1-\cdots-k_{r-2}}{k_{r-1}} \cdot x_1^{n-k_1-\cdots-k_{r-1}} x_2^{k_1} \cdots x_r^{k_{r-1}}.$$

It is immediately seen from the integral representation (2.2) that [see Eq. (1.4)]

$$\begin{aligned}
 g_n^{(\alpha_1, \dots, \alpha_r)}(x_1, \dots, x_r) &= \sum_{\substack{k_1, \dots, k_r \in \mathbf{N}_0 \\ (k_1 + \dots + k_r = n)}} \binom{\alpha_1 + k_1 - 1}{k_1} \dots \binom{\alpha_r + k_r - 1}{k_r} x_1^{k_1} \dots x_r^{k_r} \\
 &= \sum_{k_1=0}^n \sum_{k_2=0}^{n-k_1} \dots \sum_{k_{r-1}=0}^{n-k_1-\dots-k_{r-2}} \frac{(\alpha_1)_{n-k_1-\dots-k_{r-1}} (\alpha_2)_{k_1} \dots (\alpha_r)_{k_{r-1}}}{(n-k_1-\dots-k_{r-1})! k_1! \dots k_{r-2}!} \\
 (2.5) \quad &\cdot x_1^{n-k_1-\dots-k_{r-1}} x_2^{k_1} \dots x_r^{k_{r-1}},
 \end{aligned}$$

which follows also from the *second* explicit expression in (1.4).

We now define the umbral quantities  $\langle \alpha_1 x_1 \rangle, \dots, \langle \alpha_r x_r \rangle$  together with their properties given by (see, for details, [5] ; see also [14])

$$\langle \alpha_j x_j \rangle^n = (\alpha_j)_n x_j^n \quad (j = 1, \dots, r; n \in \mathbf{N}_0)$$

and the pairs of operators  $\hat{X}_j$  and  $\hat{P}_{x_j}$  ( $j = 1, \dots, r$ ) such that

$$\hat{X}_j \{ \langle \alpha_j x_j \rangle^n \} = \langle \alpha_j x_j \rangle^{n+1} \quad \text{and} \quad \hat{P}_{x_j} \{ \langle \alpha_j x_j \rangle^n \} = n \langle \alpha_j x_j \rangle^{n-1}$$

$$(j = 1, \dots, r),$$

so that, obviously,

$$\begin{aligned}
 \hat{X}_j \hat{P}_{x_j} \{ \langle \alpha_j x_j \rangle^n \} &= n \langle \alpha_j x_j \rangle^n \quad \text{and} \quad \hat{P}_{x_j} \hat{X}_j \{ \langle \alpha_j x_j \rangle^n \} = (n+1) \langle \alpha_j x_j \rangle^n \\
 (j = 1, \dots, r).
 \end{aligned}$$

We thus find that the Chan-Chyan-Srivastava polynomials  $g_n^{(\alpha_1, \dots, \alpha_r)}(x_1, \dots, x_r)$  satisfy the following operational formula:

$$\begin{aligned}
 &\left( \hat{X}_1 \hat{P}_{x_1} + \dots + \hat{X}_r \hat{P}_{x_r} \right) \left\{ g_n^{(\alpha_1, \dots, \alpha_r)}(x_1, \dots, x_r) \right\} \\
 (2.6) \quad &= n g_n^{(\alpha_1, \dots, \alpha_r)}(x_1, \dots, x_r),
 \end{aligned}$$

which, in the special case when

$$r = 2, \quad \alpha_1 = \alpha, \quad \alpha_2 = \beta, \quad x_1 = x \quad \text{and} \quad x_2 = y,$$

would provide the *corrected* version of a result stated by Dattoli *et al.* [6, p. 182, Eq. (10)].

It is also clear that the umbral image of the generating function (1.3) is given by

$$(2.7) \quad e^{[\langle \alpha_1 x_1 \rangle + \dots + \langle \alpha_r x_r \rangle]z} = \sum_{n=0}^{\infty} g_n^{(\alpha_1, \dots, \alpha_r)}(x_1, \dots, x_r) z^n.$$

In our next section (Section 3), we will show that the above notion and formalism of *monoumbrality* (that is, *monomiality* together with *umbrality*) can be extended to a large family of polynomials as well as of functions and that this principle of monoumbrality will provide a powerful tool for simplifying calculations (see also [5], [6] and [7]).

### 3. Applications of the Principle of Monoumbrality

In the preceding section, we have noted that the umbral image of the generating function (1.3) of the Chan-Chyan-Srivastava polynomials  $g_n^{(\alpha_1, \dots, \alpha_r)}(x_1, \dots, x_r)$  is, in fact, of the exponential type given by (2.7). Here, in this section, we begin by recalling the following easily understandable decomposition rules for the umbral algebra (see [5], [6] and [7]):

$$(3.1) \quad e^{\langle \alpha_1 x_1 \rangle + \dots + \langle \alpha_r x_r \rangle} = e^{\langle \alpha_1 x_1 \rangle} \dots e^{\langle \alpha_r x_r \rangle},$$

and

$$(3.2) \quad e^{[\langle \alpha(x_1 + \dots + x_r) \rangle]} \neq e^{\langle \alpha x_1 \rangle} \dots e^{\langle \alpha x_r \rangle},$$

that is, more explicitly,

$$(3.3) \quad e^{[\langle \alpha(x_1 + \dots + x_r) \rangle]} \neq \left( \sum_{k_1=0}^{\infty} (\alpha)_{k_1} \frac{x_1^{k_1}}{k_1!} \right) \dots \left( \sum_{k_r=0}^{\infty} (\alpha)_{k_r} \frac{x_r^{k_r}}{k_r!} \right).$$

We also note that

$$(3.4) \quad \begin{aligned} e^{\langle \alpha_1 x_1 \rangle z} &= e^{\langle \alpha_1 x_1 \rangle z + \dots + \langle \alpha_r x_r \rangle z} \cdot e^{\langle -\alpha_2 x_2 \rangle z} \dots e^{\langle -\alpha_r x_r \rangle z} \\ &= e^{[\langle \alpha_1 x_1 \rangle + \dots + \langle \alpha_r x_r \rangle]z} \cdot e^{[\langle -\alpha_2 x_2 \rangle + \dots + \langle -\alpha_r x_r \rangle]z}. \end{aligned}$$

By applying this last umbral relationship (3.4) in conjunction with (2.5) and (2.7), we find that

$$\begin{aligned}
 \sum_{n=0}^{\infty} (\alpha_1)_n \frac{(x_1 z)^n}{n!} &= \left( \sum_{n=0}^{\infty} g_n^{(\alpha_1, \dots, \alpha_r)}(x_1, \dots, x_r) z^n \right) \\
 &\quad \left( \sum_{n=0}^{\infty} g_n^{(-\alpha_2, \dots, -\alpha_r)}(x_2, \dots, x_r) z^n \right) \\
 (3.5) \quad &= \sum_{n=0}^{\infty} z^n \sum_{k=0}^n g_k^{(\alpha_1, \dots, \alpha_r)}(x_1, \dots, x_r) g_{n-k}^{(-\alpha_2, \dots, -\alpha_r)}(x_2, \dots, x_r),
 \end{aligned}$$

so that, upon equating the coefficients of  $z^n$  from both sides in (3.5), we finally obtain

$$\begin{aligned}
 x_1^n &= \frac{1}{(\alpha_1)_n} \sum_{k_1=0}^n \sum_{k_2=0}^{n-k_1} \cdots \sum_{k_{r-1}=0}^{n-k_1-\cdots-k_{r-2}} k_1! \binom{n}{k_1} \binom{n-k_1}{k_2} \\
 &\quad \cdots \binom{n-k_1-\cdots-k_{r-2}}{k_{r-1}} \\
 &\quad \cdot g_{k_1}^{(\alpha_1, \dots, \alpha_r)}(x_1, \dots, x_r) \langle -\alpha_2 x_2 \rangle^{n-k_1-\cdots-k_{r-1}} \langle -\alpha_3 x_3 \rangle^{k_2} \cdots \langle -\alpha_r x_r \rangle^{k_{r-1}}, \\
 (3.6) \quad &
 \end{aligned}$$

where, just as in the preceding section,

$$\begin{aligned}
 \langle -\alpha_2 x_2 \rangle^{n-k_1-\cdots-k_{r-1}} &= \frac{\Gamma(-\alpha_2 + n - k_1 - \cdots - k_{r-1})}{\Gamma(-\alpha_2)} x_2^{n-k_1-\cdots-k_{r-1}} \\
 (3.7) \quad &= (-\alpha_2)_{n-k_1-\cdots-k_{r-1}} x_2^{n-k_1-\cdots-k_{r-1}}
 \end{aligned}$$

and

$$\begin{aligned}
 \langle -\alpha_j x_j \rangle^{k_{j-1}} &= \frac{\Gamma(-\alpha_j + k_{j-1})}{\Gamma(-\alpha_j)} x_j^{k_{j-1}} = (-\alpha_j)_{k_{j-1}} x_j^{k_{j-1}} \quad (j = 3, \dots, r). \\
 (3.8) \quad &
 \end{aligned}$$

Formula (3.6) provides the expansion of an ordinary monomial  $x_1^n$  in terms of the Chan-Chyan-Srivastava polynomials

$$g_n^{(\alpha_1, \dots, \alpha_r)}(x_1, \dots, x_r)$$

involving such umbral quantities as those specified in (3.7) and (3.8).

Next, for Erkuş-Srivastava polynomials  $\mathcal{U}_{n;\ell_1,\dots,\ell_r}^{(\alpha_1,\dots,\alpha_r)}(x_1,\dots,x_r)$  generated by (1.7), it is easily seen that

$$(3.9) \quad \begin{aligned} & \mathcal{U}_{n;\ell_1,\dots,\ell_r}^{(\alpha_1+\beta_1,\dots,\alpha_r+\beta_r)}(x_1,\dots,x_r) \\ &= \sum_{k=0}^n \mathcal{U}_{n-k;\ell_1,\dots,\ell_r}^{(\alpha_1,\dots,\alpha_r)}(x_1,\dots,x_r) \mathcal{U}_{k;\ell_1,\dots,\ell_r}^{(\beta_1,\dots,\beta_r)}(x_1,\dots,x_r), \end{aligned}$$

which, for  $\ell_j = 1$  ( $j = 1, \dots, r$ ), was given by Chan *et al.* [3, p. 147, Eq. (35)]. Moreover, the generating function (1.7) together with the integral formula (2.1) would yield the following analogue of the integral representation (2.2):

$$(3.10) \quad \begin{aligned} & \mathcal{U}_{n;\ell_1,\dots,\ell_r}^{(\alpha_1+\beta_1,\dots,\alpha_r+\beta_r)}(x_1,\dots,x_r) = \frac{1}{\Gamma(\alpha_1)\cdots\Gamma(\alpha_r)} \int_0^\infty \cdots \int_0^\infty e^{-(\xi_1+\cdots+\xi_r)} \\ & \cdot \xi_1^{\alpha_1-1} \cdots \xi_r^{\alpha_r-1} \mathcal{Q}_n(x_1\xi_1, \dots, x_r\xi_r) d\xi_1 \cdots d\xi_r, \end{aligned}$$

where the polynomials  $\mathcal{Q}_n(x_1, \dots, x_r)$  are essentially the same as the multivariable Hermite-Kampé de Fériet polynomials given by

$$(3.11) \quad \mathcal{Q}_n(x_1, \dots, x_r) := \frac{1}{n!} \sum_{\substack{k_1, \dots, k_r \in \mathbf{N}_0 \\ (\ell_1 k_1 + \cdots + \ell_r k_r = n)}} \binom{n}{k_1, \dots, k_r} x_1^{k_1} \cdots x_r^{k_r}$$

in terms of the multinomial coefficients involved also in (2.3).

If we now make use of the principle of monoumbrality as detailed above, we can show similarly that

$$(3.12) \quad \begin{aligned} x_1^n &= \frac{n!}{(\alpha_1)_n} \sum_{k_1=0}^{\ell_1 n} \sum_{\substack{k_2, \dots, k_r \in \mathbf{N}_0 \\ (\ell_2 k_2 + \cdots + \ell_r k_r = \ell_1 n - k_1)}} \mathcal{U}_{k_1;\ell_1,\dots,\ell_r}^{(\alpha_1,\dots,\alpha_r)}(x_1, \dots, x_r) \\ & \cdot \frac{\langle -\alpha_2 x_2 \rangle^{k_2}}{k_2!} \cdots \frac{\langle -\alpha_r x_r \rangle^{k_r}}{k_r!}, \end{aligned}$$

which, in the special case when  $\ell_j = 1$  ( $j = 1, \dots, r$ ), the monounbrl expansion given by (3.6). Moreover, if we simply assume that  $\ell_r = 1$ , we can easily rewrite the monounbrl expansion (3.12) as follows:

$$(3.13) \quad x_1^n = \frac{n!}{(\alpha_1)_n} \sum_{k_1=0}^{\ell_1 n} \sum_{k_2=0}^{\left\lfloor \frac{\ell_1 n - k_1}{\ell_2} \right\rfloor} \cdots \sum_{k_{r-1}=0}^{\left\lfloor \frac{\ell_1 n - k_1 - \ell_2 k_2 - \cdots - \ell_{r-2} k_{r-2}}{\ell_{r-1}} \right\rfloor} \cdot \mathcal{U}_{k_1; \ell_1, \dots, \ell_{r-1}, 1}^{(\alpha_1, \dots, \alpha_r)}(x_1, \dots, x_r) \\ \cdot \frac{\langle -\alpha_2 x_2 \rangle^{\ell_1 n - k_1 - \ell_2 k_2 - \cdots - \ell_{r-1} k_{r-1}}}{(\ell_1 n - k_1 - \ell_2 k_2 - \cdots - \ell_{r-1} k_{r-1})!} \frac{\langle -\alpha_3 x_3 \rangle^{k_2}}{k_2!} \cdots \frac{\langle -\alpha_r x_r \rangle^{k_{r-1}}}{k_{r-1}!}$$

or, equivalently,

$$(3.14) \quad x_1^n = \frac{n!}{(\alpha_1)_n} \sum_{k_1=0}^{\ell_1 n} \sum_{k_2=0}^{\left\lfloor \frac{\ell_1 n - k_1}{\ell_2} \right\rfloor} \cdots \sum_{k_{r-1}=0}^{\left\lfloor \frac{\ell_1 n - k_1 - \ell_2 k_2 - \cdots - \ell_{r-2} k_{r-2}}{\ell_{r-1}} \right\rfloor} \cdot \mathcal{U}_{k_1; \ell_1, \dots, \ell_{r-1}, 1}^{(\alpha_1, \dots, \alpha_r)}(x_1, \dots, x_r) \\ \cdot \frac{k_1! (\ell_2 k_2)! \cdots (\ell_{r-1} k_{r-1})!}{k_2! \cdots k_{r-1}!} \binom{\ell_1 n}{k_1} \binom{\ell_1 n - k_1}{\ell_2 k_2} \\ \cdots \binom{\ell_1 n - k_1 - \ell_2 k_2 - \cdots - \ell_{r-2} k_{r-2}}{\ell_{r-1} k_{r-1}} \\ \cdot \langle -\alpha_2 x_2 \rangle^{\ell_1 n - k_1 - \ell_2 k_2 - \cdots - \ell_{r-1} k_{r-1}} \langle -\alpha_3 x_3 \rangle^{k_2} \cdots \langle -\alpha_r x_r \rangle^{k_{r-1}}$$

in terms of the notations and conventions given by (3.7) and (3.8). Indeed, when we *further* set  $\ell_j = 1$  ( $j = 1, \dots, r-1$ ), this last result (3.14) would correspond *precisely* to the monounbrl expansion (3.6).

Finally, upon setting  $\ell_j = j$  ( $j = 1, \dots, r$ ) in our general result (3.12), we are led at once to the following monounbrl expansion for the multivariable Lagrange-Hermite polynomials  $h_n^{(\alpha_1, \dots, \alpha_r)}(x_1, \dots, x_r)$  in (1.6), which were studied by (for example) Altın and Erkuş [1] :

$$x_1^n = \frac{n!}{(\alpha_1)_n} \sum_{k_1=0}^n \sum_{\substack{k_2, \dots, k_r \in \mathbf{N}_0 \\ (2k_2 + \cdots + rk_r = n - k_1)}} h_{k_1}^{(\alpha_1, \dots, \alpha_r)}(x_1, \dots, x_r)$$

$$(3.15) \quad \cdot \frac{\langle -\alpha_2 x_2 \rangle^{k_2}}{k_2!} \dots \frac{\langle -\alpha_r x_r \rangle^{k_r}}{k_r!}.$$

#### 4. Concluding Remarks and Observations

In this work, we have investigated the umbral calculus presentations of the Chan-Chyan-Srivastava polynomials  $g_n^{(\alpha_1, \dots, \alpha_r)}(x_1, \dots, x_r)$  generated by (1.3) and also of the substantially more general Erkuş-Srivastava polynomials  $\mathcal{U}_{n; \ell_1, \dots, \ell_r}^{(\alpha_1, \dots, \alpha_r)}(x_1, \dots, x_r)$  generated by (1.7). One of our main monoumbral expansions asserted by (3.12) has been shown to yield the corresponding monoumbral expansion (3.15) for the multivariable Lagrange-Hermite polynomials  $h_n^{(\alpha_1, \dots, \alpha_r)}(x_1, \dots, x_r)$  in (1.6), which were studied by (for example) Altın and Erkuş [1].

We need hardly emphasize upon the fact that the notion and formalism of monoumbrality can be extended to a large family of polynomials as well as of functions and that the underlying principle of monoumbrality would provide a powerful tool for simplifying calculations.

We conclude our present investigation by remarking further that, in the special case when  $\ell_1 = 1$ , we find from (1.8) that

$$\begin{aligned} \mathcal{U}_{n; 1, \ell_2, \dots, \ell_r}^{(\alpha_1, \dots, \alpha_r)}(x_1, \dots, x_r) &= \sum_{\substack{k_1, \dots, k_r \in \mathbf{N}_0 \\ (k_1 + \ell_2 k_2 + \dots + \ell_r k_r = n)}} \frac{\langle \alpha_1 x_1 \rangle^{k_1}}{k_1!} \dots \frac{\langle \alpha_r x_r \rangle^{k_r}}{k_r!} \\ &= \sum_{k_2, \dots, k_r=0}^{l_2 k_2 + \dots + l_r k_r \leq n} \frac{\langle \alpha_1 x_1 \rangle^{n - \ell_2 k_2 - \dots - \ell_r k_r}}{(n - \ell_2 k_2 - \dots - \ell_r k_r)!} \frac{\langle \alpha_2 x_2 \rangle^{k_2}}{k_2!} \dots \frac{\langle \alpha_r x_r \rangle^{k_r}}{k_r!} \\ &= \sum_{k_1=0}^{\left\lfloor \frac{n}{\ell_2} \right\rfloor} \sum_{k_2=0}^{\left\lfloor \frac{n - \ell_2 k_1}{\ell_3} \right\rfloor} \dots \sum_{k_{r-1}=0}^{\left\lfloor \frac{n - \ell_2 k_1 - \dots - \ell_{r-1} k_{r-2}}{\ell_r} \right\rfloor} \\ (4.1) \quad &\cdot \frac{\langle \alpha_1 x_1 \rangle^{n - \ell_2 k_1 - \dots - \ell_r k_{r-1}}}{(n - \ell_2 k_1 - \dots - \ell_r k_{r-1})!} \frac{\langle \alpha_2 x_2 \rangle^{k_1}}{k_1!} \dots \frac{\langle \alpha_r x_r \rangle^{k_{r-1}}}{k_{r-1}!}, \end{aligned}$$

which readily yields the following generalization of the operational formula

(2.6):

$$\left( \hat{X}_1 \hat{P}_{x_1} + \ell_2 \hat{X}_2 \hat{P}_{x_2} + \dots + \ell_r \hat{X}_r \hat{P}_{x_r} \right) \left\{ \mathcal{U}_{n; 1, \ell_2, \dots, \ell_r}^{(\alpha_1, \dots, \alpha_r)}(x_1, \dots, x_r) \right\}$$

$$(4.2) \quad = n\mathcal{U}_{n;1,\ell_2,\dots,\ell_r}^{(\alpha_1,\dots,\alpha_r)}(x_1,\dots,x_r).$$

In particular, if we set  $\ell_j = j$  ( $j = 2, \dots, r$ ) in this last result (4.2), we immediately obtain

$$(4.3) \quad \begin{aligned} & \left( \hat{X}_1 \hat{P}_{x_1} + 2\hat{X}_2 \hat{P}_{x_2} + \dots + r\hat{X}_r \hat{P}_{x_r} \right) \left\{ h_n^{(\alpha_1,\dots,\alpha_r)}(x_1,\dots,x_r) \right\} \\ & = nh_n^{(\alpha_1,\dots,\alpha_r)}(x_1,\dots,x_r) \end{aligned}$$

for the multivariable Lagrange-Hermite polynomials  $h_n^{(\alpha_1,\dots,\alpha_r)}(x_1,\dots,x_r)$  in (1.6), which were studied by (for example) Altın and Erkuş [1]. Indeed, in its *further* special case when

$$r = 2, \quad \alpha_1 = \alpha, \quad \alpha_2 = \beta, \quad x_1 = x \quad \text{and} \quad x_2 = y,$$

the operational formula (4.3) would provide the *corrected* version of another result stated by Dattoli *et al.* [6, p. 184, Eq. (22)].

## References

- [1] A. Altın and E. Erkuş, On a multivariable extension of the Lagrange-Hermite polynomials, *Integral Transforms Spec. Funct.* **17**, pp. 239–244, (2006).
- [2] L. C. Andrews, *Special Functions for Engineers and Applied Mathematicians*, Macmillan Company, New York, (1985).
- [3] W.-C. C. Chan, C.-J. Chyan and H. M. Srivastava, The Lagrange polynomials in several variables, *Integral Transforms Spec. Funct.* **12**, pp. 139–148, (2001).
- [4] K.-Y. Chen, S.-J. Liu and H. M. Srivastava, Some new results for the Lagrange polynomials in several variables, *ANZIAM J.* **49**, pp. 243–258, (2007).
- [5] G. Dattoil, P. E. Ricci and C. Cesarano, Monumbral polynomials and the associated formalism, *Integral Transforms Spec. Funct.* **13**, pp. 155–162, (2002).

- [6] G. Dattoil, P. E. Ricci and C. Cesarano, The Lagrange polynomials, the associated generalizations, and the umbral calculus, *Integral Transforms Spec. Funct.* **14**, pp. 181–186, (2003).
- [7] G. Dattoil, P. E. Ricci and C. Cesarano, Beyond the monomiality: The monumbrality principle, *J. Comput. Anal. Appl.* **6**, pp. 77–83, (2004).
- [8] A. Erdélyi, W. Magnus, F. Oberhettinger and F. G. Tricomi, *Higher Transcendental Functions*, Vol. **III**, McGraw-Hill Book Company, New York, Toronto and London, (1955).
- [9] E. Erkuş and H. M. Srivastava, A unified presentation of some families of multivariable polynomials, *Integral Transforms Spec. Funct.* **17**, pp. 267–273, (2006).
- [10] M. A. Khan and A. K. Shukla, On Lagrange’s polynomials of three variables, *Proyecciones J. Math.* **17**, pp. 227–235, (1998).
- [11] S.-J. Liu, Bilateral generating functions for the Lagrange polynomials and the Lauricella functions, *Integral Transforms Spec. Funct.* **20**, pp. 519–527, (2009).
- [12] S.-J. Liu, C.-J. Chyan, H.-C. Lu and H. M. Srivastava, Bilateral generating functions for the Chan-Chyan-Srivastava polynomials and the generalized Lauricella functions, *Integral Transforms Spec. Funct.* **23**, pp. 539–549, (2012).
- [13] S.-J. Liu, S.-D. Lin, H. M. Srivastava and M.-M. Wong, Bilateral generating functions for the Erkuş-Srivastava polynomials and the generalized Lauricella functions, *Appl. Math. Comput.* **218**, pp. 7685–7693, (2012).
- [14] S. Roman, *The Umbral Calculus*, Academic Press, New York and London, (1984).
- [15] H. M. Srivastava and H. L. Manocha, *A Treatise on Generating Functions*, Halsted Press (Ellis Horwood Limited, Chichester), John Wiley and Sons, New York, Chichester, Brisbane and Toronto, (1984).

**H. M. Srivastava**

Department of Mathematics and Statistics,  
University of Victoria,  
Victoria,  
British Columbia V8W 3R4,  
Canada  
e-mail : harimsri@math.uvic.ca

**K. S. Nisar**

Department of Mathematics,  
College of Engineering,  
Salman Bin Abdul-Aziz University,  
Wadi Dawaser,  
Kingdom of Saudi Arabia  
e-mail : ksnisar1@gmail.com

and

**Mumtaz Ahmad Khan**

Department of Applied Mathematics,  
Faculty of Engineering and Technology,  
Aligarh Muslim University,  
Aligarh 202002, Uttar Pradesh,  
India  
e-mail : mumtaz\_ahmad\_khan\_2008@yahoo.com