

SCHUR RING AND QUASI-SIMPLE MODULES

PEDRO DOMÍNGUEZ-WADE
UNIVERSIDAD DE MATANZAS, CUBA
Received : November 2008. Accepted : April 2009

Abstract

Let R be a ring of algebraic integers of an algebraic number field F and let $G \leq GL_n(R)$ be a finite group. In this paper we show that the R -span of G is just the matrix ring $M_n(R)$ of the $n \times n$ -matrices over R if and only if $G/O_{p_i}(G)$ is absolutely simple for all $p_i \in \pi$, where π is the set of the positive prime divisors of $|G|$ and $O_{p_i}(G)$ is the largest normal p_i -subgroup.

Subclass : *Primary 20C20 ; Secondary 19A22.*

Keywords : *Schur ring, π -globally simple.*

1. Introduction.

Let F be an algebraic number field with ring of algebraic integers R and let $\pi = \{p_1, \dots, p_t\}$ be a set of positive prime numbers. Assume that the I_i are maximal ideals of R such that $p_i \in I_i (i = 1, \dots, t)$. Set $L_\pi = \{\frac{a}{b} \mid a, b \in R, b \notin I_i, i = 1, \dots, t\}$. Then R_π denotes a localization of R at L_π . Thus R_π is a principal ideal having quotient field of characteristic zero and containing a unique prime ideal I_i such that $p_i \in I_i, i = 1, \dots, n$. We denote the Jacobson radical of R_π be $J(R_\pi)$. Therefore the residue ring $K_m = R_\pi/J(R_\pi)$ is a semi-simple ring of characteristic m . We may write

$$(1.1) \quad K_m = \bigoplus_{i=1}^t k_i$$

where the k_i are fields of characteristic $p_i (i = 1, \dots, t)$.

If G is a finite group then we obtain

$$(1.2) \quad K_m G = \bigoplus_{i=1}^t k_i G.$$

by (1.1).

From (1.2) it follows that

$$(1.3) \quad 1 = f_1 + \dots + f_t$$

where the f_i are orthogonal central idempotents in $K_m G$.

Therefore

$$K_m G = \bigoplus_{i=1}^t k_i G \\ = \bigoplus_{i=1}^t K_m G f_i \text{ with } k_i G = K_m G f_i.$$

Now, R_π is a Hausdorff space in its $J(R_\pi)$ -topology, i. e., that

$$\bigcap_{j=1}^{\infty} J(R_\pi)^j = (0).$$

Therefore the $J(R_\pi)$ -adic completion $\hat{R}_\pi$ of R_π is a complete semi-local ring such that

$$K_m = R_\pi/J(R_\pi) = \hat{R}_\pi/J(\hat{R}_\pi)$$

and

$$(1.4) \quad \hat{R}_\pi = R_1 \oplus \dots \oplus R_t$$

where the R_i are complete local rings such that $R_i/J(R_i) \cong k_i$.

Observe that R_i is isomorphic to $\hat{R}_{v_{p_i}}$, where $\hat{R}_{v_{p_i}}$ is a complete valuation ring corresponding to the discrete valuation v_{p_i} associated to the maximal ideal I_i of R . Thus, we may write

$$(1.5) \quad \hat{R}_\pi \cong \hat{R}_{v_{p_1}} \oplus \cdots \oplus \hat{R}_{v_{p_t}}$$

by (1.4).

Therefore

$$(1.6) \quad \hat{R}_\pi G \cong \hat{R}_{v_{p_1}} G \oplus \cdots \oplus \hat{R}_{v_{p_t}} G.$$

From (1.6) we obtain

$$(1.7) \quad 1 = \hat{f}_1 + \cdots + \hat{f}_n$$

where the $\hat{f}_i$ are orthogonal central idempotents in $\hat{R}_\pi G$.

Let π_l be any set of positive prime numbers. Assume that R_{π_l} is the localization of R at L_{π_l} . Then we have

$$(1.8) \quad \bigcap_{l=1}^{\infty} R_{\pi_l} = R$$

In the study of the Schur ring $M_n(R)$ of a commutative ring R the main problem is to find a finite group $G \leq GL_n(R)$ such that the R -span of G coincides with the matrix ring $M_n(R)$ of the $n \times n$ -matrices on K . The more precise question, in general sense, which Azumaya algebras over R are obtainable as an epimorphic image of the group-ring RG for some finite group G .

1.1. Notations and Definitions.

Throughout the paper F denote an algebraic number field and R denote the ring of algebraic integers of F . Moreover, K_m is semi-simple ring of characteristic m with maximal ideals K_i and residue fields $k_i = K_m/K_i$ of characteristic p_i . For an maximal ideal I of R we denote by ϕ the I -adic valuation on F . Here R_{v_p} denote the valuation ring of v_p and $\hat{R}_{v_p}$ denote the complete valuation ring corresponding to the discrete valuation v_p . Let $M_n(R)$ stand for the ring of $(n \times n)$ -matrices over R . We write $GL_n(R)$ for the multiplicative group of the invertible elements of $M_n(R)$. For a finite subgroup G of $GL_n(R)$ we let $\langle G \rangle_R$ be the R -span of G in $M_n(R)$. Let π be a set of natural primes. We denote the fields of rational and complex numbers by $\mathbf{Q}$ and $\mathbf{C}$, respectively.

2. Preliminary Results.

Let R be a commutative ring and let $G \leq GL_n(R)$ be a finite group. Then the matrix ring $M_n(R)$ is called Schur ring if $\langle G \rangle_R = M_n(R)$.

Lemma 2.0.1. *Let k be a field of characteristic p and let $G \leq GL_n(k)$ be a finite group. Assume that V is the kG -module corresponding to G . Then G is absolutely simple if and only if $\langle G \rangle_k = M_n(k)$.*

Proof. If G is absolutely simple then by Burnside's theorem the assertion follows. Conversely, let $\dot{G}$ be a finite group with a representation $\varphi : \dot{G} \rightarrow GL_n(k)$ such that $\varphi(\dot{G}) = G$. Consider the surjection of k -algebras $\psi : k\dot{G} \rightarrow \langle G \rangle_k$, where $\psi(\dot{G}) = \varphi(\dot{G})$. Therefore $k\dot{G}/\ker \psi \cong M_n(k)$. Since $J(k\dot{G}) \subseteq \ker \psi$ it follows that the matrix algebra summand of $k\dot{G}/J(k\dot{G})$ corresponding to V is $M_n(k)$, so the result follows. $\square$

Let K_m be a semi-simple ring of characteristic m and let $G \leq GL_n(K_m)$ be a finite group. Assume that V is the $K_m G$ -module corresponding to G . Then G is called π -quasi-simple if each direct summand $V f_i$ is an absolutely simple $k_i G$ -module.

Lemma 2.0.2. *Let K_m be a semi-simple ring of characteristic m and let $G \leq GL_n(K_m)$ be a finite group. Assume that V is the $K_m G$ -module corresponding to G . Then $\langle G \rangle_{K_m} = M_n(K_m)$ if and only if G is π -quasi-simple.*

Proof. We have

$$\begin{aligned} \langle G \rangle_{K_m} &= \langle G \rangle_{k_1} \oplus \cdots \oplus \langle G \rangle_{k_t} \\ &= M_n(K_m) \\ &= M_n(k_1) \oplus \cdots \oplus M_n(k_t) \end{aligned}$$

Therefore $\langle G \rangle_{k_i} = M_n(k_i)$. Thus applying the last lemma the result follows. Conversely, applying again the lemma (2.0.1) we deduce that

$$\langle G \rangle_{k_i} = M_n(k_i)$$

for all i . Thus we may write

$$\begin{aligned} \langle G \rangle_{K_m} &= \langle G \rangle_{k_1} \oplus \cdots \oplus \langle G \rangle_{k_t} \\ &= M_n(k_1) \oplus \cdots \oplus M_n(k_t) \\ &= M_n(K_m) \end{aligned}$$

So we are done.

□

Proposition 2.0.3. *Let $G \leq GL_n(\mathbf{C})$ be a finite group. Assume that F is a finite extension of $\mathbf{Q}$ such that $G \leq GL_n(F)$. If R is the ring of integers of F then $G \leq GL_n(R_\pi)$, being R_π a localization of R at L_π .*

Proof. Let $\hat{R}_\pi$ be the completion of R_π . It is well known that $G \leq GL_n(\hat{R}_{v_{p_l}})(l = 1, \dots, t)$, where $\hat{R}_{v_{p_l}}$ is a complete valuation ring. Let $\hat{U}_i$ be the $\hat{R}_{v_{p_l}}G$ -module corresponding to G . From (1.5) we deduce that there is an $\hat{R}_\pi G$ -module minimal $\hat{R}_\pi$ -free $\hat{U} = \hat{U}_1 \oplus \dots \oplus \hat{U}_t$, so $G \leq GL_n(\hat{R}_\pi)$. Since $\hat{U}$ is a complete minimal $\hat{R}_\pi$ -free module it follows that $\hat{U} = \hat{R}_\pi U$, being U an $R_\pi G$ -module, which is a Hausdorff space for its $J(R_\pi)$ -topology. So we are done. □

From (1.8) and the last proposition one can deduce an “Hasse Principle” for $\langle G \rangle_R$ to coincide with $M_n(R)$: this is the case if and only if $\langle G \rangle_{R_\pi} = M_n(R_\pi)$ for every set of positive prime numbers π .

Let $G \leq GL_n(\mathbf{C})$ be a finite group and let $U_{\mathcal{C}}$ be the $\mathbf{C}G$ -module corresponding to G . Then there exists a finite extension F of $\mathbf{Q}$ with FG -module U_F conjugate to $U_{\mathcal{C}}$. According to the proposition (2.0.3) there is an $R_\pi G$ -module U which is also conjugate to $U_{\mathcal{C}}$. Let $\hat{R}_\pi$ be the $J(R_\pi)$ -adic completion of R_π . We know that there exists an $\hat{R}_\pi G$ -module $\hat{U}$ which is the completion of U . From (1.7) it follows that

$$\hat{U} = \hat{U}\hat{f}_1 \oplus \dots \oplus \hat{U}\hat{f}_t$$

where the $\hat{U}\hat{f}_i$ are $R_{v_{p_i}}G$ -modules. The $K_m G$ -module $\overline{U}_{K_m} = \hat{U}/J(R_\pi)\hat{U} = \hat{U}\hat{f}_1/J(R_{v_{p_1}})\hat{U}\hat{f}_1 \oplus \dots \oplus \hat{U}\hat{f}_t/J(R_{v_{p_t}})\hat{U}\hat{f}_t$

is called reduction of U modulo π .

The natural projection $GL_n(\hat{R}_\pi) \rightarrow GL_n(K_m)$ induces the homomorphism $\tau : G \rightarrow GL_n(K_m)$, where $\tau(G) = \overline{G}$. Furthermore the homomorphism $GL_n(\hat{R}_{v_{p_i}}) \rightarrow GL_n(k_i)$ induces the homomorphism $\tau_i : G \rightarrow GL_n(k_i)$ with $\tau_i(G) = \overline{G}_i$

Observe that if $\langle G \rangle_{R_{\pi_l}} = M_n(R_{\pi_l})$, where π_l is the set of the positive prime divisors of $|G|$, then $\langle G \rangle_{R_\pi} = M_n(R_\pi)$ for every set π of positive prime numbers. We use Nakayama’s lemma to show that that $\langle G \rangle_{R_{\pi_l}} = M_n(R_{\pi_l})$ is equivalent to $\langle \overline{G} \rangle_{K_m} = M_n(K_m)$.

3. Main Results.

Let R be a ring of algebraic integers and let $G \leq GL_n(R)$ be a finite group. Assume that π is the set of the positive prime divisors of $|G|$ and that U is the $R_\pi G$ -module corresponding to G . If the reduction of U module π is a π -quasi-simple $K_m G$ -module, then we say that G is a π -globally simple.

Lemma 3.0.4. *Let R be a ring of algebraic integers with localization R_π and let $G \leq GL_n(R_\pi)$ be a finite group. Assume that π is a set of the positive prime divisors of $|G|$. Then $\langle G \rangle_{R_\pi} = M_n(R_\pi)$ if and only if G is π -globally simple.*

Proof. From $\langle G \rangle_{R_\pi} = M_n(R_\pi)$ we deduce that $\langle \overline{G} \rangle_{K_m} = M_n(K_m)$, being $K_m = R_\pi/J(R_\pi)$ a semi-simple ring. The result follows by lemma (2.0.2). Conversely, applying again the lemma (2.0.2) we obtain $\langle \overline{G} \rangle_{K_m} = M_n(K_m)$, so the assertion follows. $\square$

Theorem 3.0.5. *Let R be a ring of algebraic integers and let $G \leq GL_n(R)$ be a finite group. Then $\langle G \rangle_R = M_n(R)$ if and only if G is π -globally simple.*

Proof. From $\langle G \rangle_R = M_n(R)$ we deduce that $\langle G \rangle_{R_\pi} = M_n(R_\pi)$. Hence we obtain $\langle \overline{G} \rangle_{K_m} = M_n(K_m)$. The result follows by lemma (2.0.2). Conversely, since

$$\langle G \rangle_{R_\pi} = M_n(R_\pi)$$

by lemma (3.0.4) we deduce that for any set of prime numbers π we obtain

$$\langle G \rangle_{R_\pi} = M_n(R_\pi).$$

So we are done. $\square$

Theorem 3.0.6. *Let $G \leq GL_n(\mathbf{C})$ be a finite group. Then G is π -globally simple if and only if $G/O_{p_i}(G)$ is absolutely simple.*

Proof. Let F be a finite extension of $\mathbf{Q}$ such that $G \leq GL_n(F)$, and let R be the ring of integers of F . Consider the homomorphism $h_i : G \rightarrow \overline{G}_i$. Since $\overline{G}_i$ is absolutely simple it follows that $\ker h_i = O_{p_i}(G)$. Conversely, since $G/O_{p_i}(G)$ is absolutely simple for all i , we deduce that G is π -quasi-simple, being π the set of positive prime divisors of the order of G . So we are done. $\square$

References

- [1] P. HUU TIEP. Globally irreducible representations of the finite symplectic group $\mathrm{Sp}_4(q)$, Comm. in Algebra 22, pp. 6439 - 6457, (1994).
- [2] P. HUU TIEP. Basic spin representations of $2\mathrm{S}_n$ and $2\mathrm{A}_n$ as globally irreducible representations, Archiv Math. 64, pp. 103 - 112, (1995).
- [3] P. HUU TIEP. Weil representations as globally irreducible representations, Math. Nachr. 184, pp. 313 - 327, (1997).
- [4] P. HUU TIEP. Globally irreducible representations of finite groups and integral lattices, Geomet. Dedicata 64, pp. 85 - 123, (1997).
- [5] P. SPIGA AND Q. WANG An answer to Hirasaka and Muzychuk : Every p -Schur ring over C_p^3 is Schurian, Discrete Mathematics 308, pp. 1760-1763, (2008).
- [6] A. E. ZALESSKII AND F. VAN OYSTAEYEN Finite Groups over Arithmetical Rings and Globally Irreducible Representations, J. Algebra 215, pp. 418-436, (1999).

Pedro Domínguez Wade

Department of Mathematics

Matanzas University

Carretera Matanzas

Varadero Km. 3

Cuba

e-mail : pedro.dominguez@umcc.cu