

Proyecciones

Vol. 28, N° 1, pp. 21–26, Mayo 2009.

Universidad Católica del Norte

Antofagasta - Chile

DOI: 10.4067/S0716-09172009000100002

## ARENS REGULARITY OF SOME BILINEAR MAPS

*M. ESHAGHI GORDJI*  
*SEM NAN UNIVERSITY, IRÁN*

*Received : February 2008. Accepted : November 2008*

### Abstract

*Let  $H$  be a Hilbert space. we show that the following statements are equivalent: (a)  $B(H)$  is finite dimension, (b) every left Banach module action  $l : B(H) \times H \longrightarrow H$ , is Arens regular (c) every bilinear map  $f : B(H)^* \longrightarrow B(H)$  is Arens regular. Indeed we show that a Banach space  $X$  is reflexive if and only if every bilinear map  $f : X^* \times X \longrightarrow X^*$  is Arens regular.*

**Subjclass [2000] :** *Primary 46H25, 16E40.*

**Keywords :** *Banach algebra, Bilinear map, Arens products*

## 1. Introduction

The regularity of bilinear maps on norm spaces, was introduced by Arens in 1951. Let  $X$ ,  $Y$  and  $Z$  be normed spaces and let  $f : X \times Y \longrightarrow Z$  be a continuous bilinear map, then  $f^* : Z^* \times X \longrightarrow Y^*$  (the transpose of  $f$ ) is defined by

$$\langle f^*(z^*, x), y \rangle = \langle z^*, f(x, y) \rangle \quad (z^* \in Z^*, x \in X, y \in Y).$$

$f^*$  is a continuous bilinear map.

Set  $f^{**} = (f^*)^*$  and  $f^{***} = (f^{**})^*, \dots$ . Then  $f^{***} : X^{**} \times Y^{**} \longrightarrow Z^{**}$  is the unique extension of  $f$  such that  $f^{***}(\cdot, y'') : X^{**} \longrightarrow Z^{**}$  is *weak\** – *weak\** continuous for every  $y'' \in Y^{**}$ . Let  $f^r : Y \times X \longrightarrow Z$  defined by  $f^r(y, x) = f(x, y)$ , ( $x \in X, y \in Y$ ), then  $f^r$  is a continuous bilinear map.  $f$  is called Arens regular whenever  $f^{***} = f^{r***r}$ . It is easy to show that  $f$  is Arens regular if and only if  $f^{***}(x'', \cdot) : Y^{**} \longrightarrow Z^{**}$  is *weak\** – *weak\** continuous for every  $x'' \in X^{**}$ . For further details we refer the reader to [A], [E-F], [D-R-V], and [M-Y]. Let  $\mathcal{A}$  be a Banach algebra and let  $\pi : \mathcal{A} \times \mathcal{A} \longrightarrow \mathcal{A}$  be the product of  $\mathcal{A}$ . Then the second dual space  $\mathcal{A}^{**}$  of  $\mathcal{A}$  is Banach algebra with both products  $\pi^{***}$  and  $\pi^{r***r}$ .  $\mathcal{A}$  is called Arens regular if  $\pi$  is Arens regular. See [A], [D-H] and [F-S]. An element  $f \in \mathcal{A}^*$  is called weakly almost periodic if the maps

$$a \mapsto a.f, \quad a \mapsto f.a, \quad \mathcal{A} \longrightarrow \mathcal{A}^*$$

are weakly compact. The collection of weakly almost periodic elements in  $\mathcal{A}^*$  is denoted by  $Wap(\mathcal{A}^*)$ .

**1.Theorem.** Let  $X$  be a Banach space and let  $l : B(X) \times X \longrightarrow X$  defined by  $l(T, x) = T(x)$ , ( $T \in B(X), x \in X$ ). If  $l$  is Arens regular, then  $B(X)^{**}$  is isomorphic to a subalgebra of  $B(X^*)$ .

**Proof:** Let  $\pi : B(X) \times B(X) \longrightarrow B(X)$  be the product of  $B(X)$ . Then for every  $e \in X$ ,  $e' \in X^*$ , and  $T, S \in B(X)$ , we have

$$\begin{aligned} \langle l^*(e', \pi(T, S)), e \rangle &= \langle e', l(T, S(e)) \rangle \\ &= \langle l^*(e', T), l(S, e) \rangle \\ &= \langle l^*(l^*(e', T)S), e \rangle. \end{aligned}$$

This means that

$$l^*(e', \pi(T, S)) = l^*(L^*(e', T), S) \quad (1).$$

By applying (1), for every  $e'' \in X^{**}$ , we have

$$\begin{aligned} \langle \pi^*(l^{**}(e'', e'), T), S \rangle &= \langle l^{**}(e'', e'), \pi(T, S) \rangle \\ &= \langle e'', l^*(e', \pi(T, S)) \rangle \\ &= \langle e'', l^*(L^*(e', T), S) \rangle \\ &= \langle l^{**}(e'', L^*(e', T), S) \rangle. \end{aligned}$$

Thus we have

$$\pi^*(l^{**}(e'', e'), T) = l^{**}(e'', L^*(e', T)) \quad (2).$$

Let  $B \in B(X)^{**}$ , then by (2), we have

$$\begin{aligned} \langle \pi^{**}(B, l^{**}(e'', e'), T) \rangle &= \langle B, \pi^*(l^*(e'', e'), T) \rangle \\ &= \langle B, l^{**}(e'', L^*(e', T)) \rangle \\ &= \langle l^{***}(B, e''), L^*(e', T) \rangle \\ &= \langle l^{**}(l^{***}(B, e''), e'), T \rangle. \end{aligned}$$

This means that

$$\pi^{**}(B, l^{**}(e'', e')) = l^{**}(l^{***}(B, e''), e') \quad (3).$$

Let now  $A \in B(X)^{**}$ . By (3), we have

$$\begin{aligned} \langle l^{***}\pi^{***}(A, B), e'' \rangle &= \langle \pi^{***}(A, B), l^{**}(e'', e') \rangle \\ &= \langle A, \pi^{**}(B, l^{**}(e'', e')) \rangle \\ &= \langle A, l^{**}(l^{***}(B, e''), e') \rangle. \end{aligned}$$

Therefore

$$l^{***}(\pi^{***}(A, B), e'') = l^{***}(A, l^{***}(B, e'')).$$

This means that the mapping  $f : B(X)^{**} \longrightarrow B(X^{**})$ , defined by  $f(A) = l^{***}(A, \cdot)$  ( $A \in B(X)^{**}$ ), is a Banach algebras homomorphism. On the other hand  $l$  is Arens regular, then  $f(A) : X^{**} \longrightarrow X^{**}$  is *weak\** – *weak\** continuous for every  $A \in B(X)^{**}$ . Let now

$$B_{w^*}(X^{**}) := \{U \in B(X^{**}) \mid U : X^{**} \rightarrow X^{**} \text{ is } \textit{weak}^* - \textit{weak}^* \text{ continuous} \},$$

and let  $\phi : B(X^*) \longrightarrow B_{w^*}(X^{**})$  defined by  $\phi(T) = T^*$  ( $T \in B(X^*)$ ). Then  $\phi^{-1} \circ f : B(X)^{**} \longrightarrow B(X^*)$  is a injective Banach algebras anti homomorphism.

**2.Theorem.** Let  $\mathcal{A}$  be a Banach algebra and let  $L$  be a reflexive left Banach  $\mathcal{A}$  module with module action  $g : \mathcal{A} \times \mathcal{L} \longrightarrow \mathcal{L}$ . Then  $g^{**}(L^{**} \times L^*) \subseteq Wap(\mathcal{A}^*)$ .

**Proof:** Let  $x'' \in L^{**}$ ,  $x' \in X^*$ , and let  $\{a_n\}$  and  $\{b_n\}$  be sequences in  $\mathcal{A}$ , in which the limits:

$$\lim_m \lim_n \langle g^{**}(x'', x'), a_m a_n \rangle, \quad \text{and} \quad \lim_n \lim_m \langle g^{**}(x'', x'), a_m a_n \rangle,$$

are both exist. Then we have

$$\begin{aligned} \lim_m \lim_n \langle g^{**}(x'', x'), a_m b_n \rangle &= \lim_m \lim_n \langle x'', g^*(x', a_m b_n) \rangle \\ &= \lim_m \lim_n \langle x'', g^*(g^*(x', a_m), b_n) \rangle \\ &= \lim_m \lim_n \langle g^{**}(x'', g^*(x', a_m)), b_n \rangle \\ &= \lim_m \lim_n \langle g^{***}(\widehat{b_n}, x''), g^*(x', a_m) \rangle \\ &= \lim_n \lim_m \langle g^{***}(\widehat{b_n}, x''), g^*(x', a_m) \rangle \quad (L \text{ is reflexive}) \\ &= \lim_n \lim_m \langle g^{**}(x'', g^*(x', a_m)), b_n \rangle \\ &= \lim_n \lim_m \langle x'', g^*(g^*(x', a_m), b_n) \rangle \\ &= \lim_n \lim_m \langle x'', g^*(x', a_m b_n) \rangle \\ &= \lim_m \lim_n \langle g^{**}(x'', x'), a_m b_n \rangle. \end{aligned}$$

This means that  $g^{**}(x'', x') \in Wap(\mathcal{A}^*)$  [Pa, Theorem 1.4.11].

**3.Corollary.** Let  $\mathcal{A}, \mathcal{L}$  and  $g$  are as above. If  $g^{**}(L^{**} \times L^*) = \mathcal{A}^*$ , then  $\mathcal{A}$  is Arens regular.

**4.Corollary [D].** Let  $X$  be a Banach space and let  $L$  be a reflexive left Banach  $B(X)$  module. We define the map  $\phi : L \widehat{\otimes} L^* \longrightarrow B(X)^*$  by

$$\langle \phi(f \otimes \mu), T \rangle = \langle \mu, T.f \rangle \quad (f \otimes \mu \in L \widehat{\otimes} L^*, T \in B(X)).$$

If  $\phi$  is surjective then  $B(X)$  is Arens regular.

**5.Theorem.** Let  $X$  be a Banach space. Then  $X$  is reflexive if and only if every bilinear map  $f : X^* \times X \longrightarrow X^*$  is Arens regular.

**Proof:** Obviously every bilinear map  $f : X^* \times X \longrightarrow X^*$  is Arens regular if  $X$  is reflexive. For the converse, let  $0 \neq x_0 \in X$ . Then by Hahn-Banach Theorem, there exists  $g \in X^*$  such that  $\langle g, x_0 \rangle = 1$ . Let  $f : X \times X \longrightarrow X^*$  defined by  $f((x, y)) = \langle g, y \rangle x$ .  $f$  is a bilinear map. Then  $f^* : X^* \times X \longrightarrow X^*$  is Arens regular. Let now  $x''' \in X^{***}$  and let  $x''_\alpha \xrightarrow{weak^*} x''$  in  $X^{**}$ , then  $f^{****}(x''', x''_\alpha) \xrightarrow{weak^*} f^{****}(x''', x'')$  in  $X^{***}$ . Thus for every  $y'' \in X^{**}$  we

have

$$\begin{aligned} \lim_{\alpha} \langle x''', f^{***}(x''_{\alpha}, y'') \rangle &= \lim_{\alpha} \langle f^{****}(x''', x''_{\alpha}, y'') \rangle \\ &= \langle f^{****}(x''', x'', y'') \rangle \\ &= \langle x''', f^{***}(x'', y'') \rangle \quad (4). \end{aligned}$$

On the other hand for each  $y'' \in X^{**}$ , there exists a net  $(y_{\beta})$  in  $Y$  such that  $\widehat{y_{\beta}} \xrightarrow{weak^*} y''$  in  $X^{**}$ . We know that  $f^{***}(\cdot, \widehat{x_0}) : X^{**} \rightarrow X^{**}$  is  $weak^* - weak^*$  continuous, then

$$f^{***}(y'', \widehat{x_0}) = w^* - \lim_{\beta} f^{***}(\widehat{y_{\beta}}, \widehat{x_0}) = w^* - \lim_{\beta} \widehat{y_{\beta}} = y'' \quad (5).$$

By (4) and (5), we have

$$\begin{aligned} \lim_{\alpha} \langle x''', x''_{\alpha} \rangle &= \lim_{\alpha} \langle x''', f^{***}(x''_{\alpha}, \widehat{x_0}) \rangle \\ &= \langle x''', f^{***}(x'', \widehat{x_0}) \rangle \\ &= \langle x''', x'' \rangle. \end{aligned}$$

This means that  $x''' : X^{**} \rightarrow C$  is  $weak^* - weak^*$  continuous. Thus  $x''' \in \widehat{X^*}$ , and  $X^*$  is reflexive. So  $X$  is reflexive.

**6. Corollary.** Let  $H$  be Hilbert space. Then the following assertions are equivalent:

- (a)  $B(H)$  is finite dimension.
- (b) The mapping  $l : B(H) \times H \rightarrow H$  defined by  $l(T, x) = T(x)$ ,  $(T \in B(H), x \in H)$ , is Arens regular.
- (c) Every bilinear map  $f : B(H)^* \times B(H) \rightarrow B(H)^*$  is Arens regular.

**Proof.** (a)  $\Leftrightarrow$  (b): Let  $B(H)$  be finite dimension, the  $B(H)$  is reflexive. Then by Theorem 1,  $l$  is Arens regular. For the converse we know that  $H \cong H^*$  as Banach spaces. Then by Theorem 1, we have  $B(H)^{**} = B(H)$ , so  $B(H)$  is finite dimension.

(a)  $\Leftrightarrow$  (c): It follows from the Theorem 5.

**Example.** Let  $H = l^1(N)$ . Then the bilinear map  $l : B(H) \times H \rightarrow H$  defined by  $l(T, x) = T(x)$ ,  $(T \in B(H), x \in H)$ , is not Arens regular.

**Acknowledgements:** The author would like to express his sincere thanks to referee for his invaluable comments. Also he would like to thank the Semnan University for its financial support.

## References

- [A] R. Arens, The adjoint of a bilinear operation, Proc. Amer. Math. Soc. 2(1951), pp. 839-848, (1951).
- [D-R-V] H. G. Dales, A. Rodriguez-Palacios and M. V. Velasco, The second transpose of a derivation, J. London Math. Soc. (2) 64, pp. 707-721, (2001).
- [D] M. Daws, Arens regularity of the algebra of operators on a Banach space. Bull. London Math. Soc. 36, No. 4, pp. 493-503, (2004).
- [D-H] J. Duncan and S. A. Hosseiniun, The second dual of Banach algebra, Proc. Roy. Soc. Edinburgh Set. A **84**, pp. 309-325, (1979).
- [E-F] M. Eshaghi Gordji and M. Filali, Arens regularity of module actions, Studia Math., Vol 181, No 3, pp. 237-254, (2007).
- [F-S] M. Filali and A. I. Singh, Recent developments on Arens regularity and ideal structure of the second dual of a group algebra and some related topological algebras. General topological algebras (Tartu, 1999), pp. 95-124, Math. Stud. (Tartu), 1, Est. Math. Soc., Tartu, 2001. MR1853838
- [M-Y] Gh. Mohajeri-Minaei, T. Yazdanpanah, Arens regularity of modules. J. Inst. Math. Comput. Sci. Math. Ser. 18, No. 3, pp. 195-197. MR2194165, (2005).
- [Pa] Theodore W. Palmer, *Banach Algebra and The General Theory of \*-Algebras Vol 1* Cambridge University Press,(1994).

**M. Eshaghi Gordji**

Department of Mathematics

Semnan University,

P. O. Box 35195-363,

Semnan,

Iran

e-mail : maj\_ess@yahoo.com ; madjid.eshaghi@gmail.com