

Proyecciones  
Vol. 26, N° 2, pp. 145-156, August 2007.  
Universidad Católica del Norte  
Antofagasta - Chile DOI: 10.4067/S0716-09172007000200001

## ON SOME PROPERTIES OF A CLASS OF POLYNOMIALS SUGGESTED BY MITTAL

*A. K. SHUKLA*

*and*

*J. C. PRAJAPATI*

*S. V. NATIONAL INSTITUTE OF TECHNOLOGY, INDIA*

*Received : September 2006. Accepted : March 2007*

### Abstract

*The object of this paper is to establish some generating relations by using operational formulae for a class of polynomials  $T_{kn}^{(\alpha+s-1)}(x)$  defined by Mittal. We have also derived finite summation formulae for (1.6) by employing operational techniques. In the end several special cases are discussed.*

**Key Words :** *Operational formulae; generating relations; finite sum formulae.*

**2000 Mathematics Subject Classification :** *33E12; 33E99; 44A45.*

## 1. Introduction

Chak [1] defined a class of polynomials as:

$$(1.1) \quad G_{n,k}^{(\alpha)}(x) = x^{-\alpha-kn+n} e^x (x^k D)^n [x^\alpha e^{-x}]$$

where  $D = \frac{d}{dx}$ ,  $k$  is constant and  $n = 0, 1, 2, \dots$ .

Chatterjea [2] studied a class of polynomials for generalized Laguerre polynomial as:

$$(1.2) \quad T_{rn}^{(\alpha)}(x, p) = \frac{1}{n!} x^{-\alpha-n-1} \exp(px^r) (x^2 D)^n [x^{\alpha+1} \exp(-px^r)].$$

Gould and Hopper [3] introduced generalized Hermite polynomials as:

$$(1.3) \quad H_n^r(x, a, p) = (-1)^n x^{-a} \exp(px^r) D^n [x^a \exp(-px^r)].$$

Singh [10] obtained generalized Truesdell polynomials by using Rodrigues formula, which is defined as:

$$(1.4) \quad T_n^{(\alpha)}(x, r, p) = x^{-\alpha} \exp(px^r) (xD)^n [x^\alpha \exp(-px^r)].$$

In 1971, Mittal [5] proved the Rodrigues formula for a class of polynomials  $T_{kn}^{(\alpha)}(x)$  as:

$$(1.5) \quad T_{kn}^{(\alpha)}(x) = \frac{1}{n!} x^{-\alpha} \exp\{p_k(x)\} D^n [x^{\alpha+n} \exp\{-p_k(x)\}]$$

where  $p_k(x)$  is a polynomial in  $x$  of degree  $k$ .

Mittal [6] also proved the following relation for (1.5)

$$(1.6) \quad T_{kn}^{(\alpha+s-1)}(x) = \frac{1}{n!} x^{-\alpha-n} \exp\{p_k(x)\} \theta^n [x^\alpha \exp\{-p_k(x)\}]$$

and an operator  $\theta \equiv x(s + xD)$ , where  $s$  is constant.

The following well-known facts are prepared for studying (1.6).

*Generalised Laguerre polynomials* (Srivastava and Manocha[12]) defined as:

$$(1.7) \quad L_n^{(\alpha)}(x) = \frac{x^{-\alpha-n-1} e^x}{n!} (x^2 D)^n [x^{\alpha+1} e^{-x}].$$

*Hermite polynomials* (Rainville [9]) defined as:

$$(1.8) \quad H_n(x) = (-1)^n \exp(x^2) D^n[\exp(-x^2)].$$

*Konhauser polynomials of first kind* (Srivastava [11]) defined as:

$$(1.9) \quad Y_n^\alpha(x; k) = \frac{x^{-kn-\alpha-1} e^x}{k^n n!} (x^{k+1} D)^n [x^{\alpha+1} e^{-x}].$$

*Konhauser polynomials of second kind* (Srivastava [11]) defined as:

$$(1.10) \quad Z_n^\alpha(x; k) = \frac{\Gamma(kn + \alpha + 1)}{n!} \sum_{j=0}^n (-1)^j \binom{n}{j} \frac{x^{kj}}{\Gamma(kj + \alpha + 1)}$$

where  $k$  is a positive integer.

Srivastava and Manocha [12] verified following result by using induction method,

$$(1.11) \quad (x^2 D)^n \{f(x)\} = x^{n+1} D^n \{x^{n-1} f(x)\}.$$

## 2. Definitions and Notations

McBride [4] defined generating function as:

Let  $G(x, t)$  be a function that can be expanded in powers of  $t$  such that

$G(x, t) = \sum_{n=0}^{\infty} c_n f_n(x) t^n$ , where  $c_n$  is a function of  $n$  that may contain the parameters of the set  $\{f_n(x)\}$ , but is independent of  $x$  and  $t$ . Then  $G(x, t)$  is called a generating function of the set  $\{f_n(x)\}$ .

**Remark:** A set of functions may have more than one generating function.

In our investigation we used the following properties of the differential operators;

$\theta \equiv x(s + xD)$  and  $\theta_1 \equiv (1 + xD)$ , where  $D \equiv \frac{d}{dx}$ , (Mittal [7], Patil and Thakare [8]) which are useful to establish linear generating relations and finite sum formulae.

$$(2.1) \quad \theta^n = x^n(s + xD)(s + 1 + xD)(s + 2 + xD) \dots (s + (n - 1) + xD)$$

$$(2.2) \quad \theta^n(x^\alpha) = (\alpha + s)_n x^{\alpha+n}$$

$$(2.3) \quad \theta^n(xuv) = x \sum_{m=0}^{\infty} \binom{n}{m} \theta^{n-m}(v) \theta_1^m(u)$$

$$(2.4) \quad e^{t\theta}(x^\alpha) = x^\alpha(1 - xt)^{-(\alpha+s)}$$

$$(2.5) \quad e^{t\theta}(xuv) = xe^{t\theta}(v)e^{t\theta_1}(u)$$

$$(2.6) \quad e^{t\theta}(x^\alpha f(x)) = x^\alpha(1 - xt)^{-(\alpha+s)} f\left[x(1 - xt)^{-1}\right]$$

$$(2.7) \quad e^{t\theta}(x^{\alpha-n} f(x)) = x^\alpha(1 + t)^{-1+(\alpha+s)} f\left[x(1 + t)\right]$$

$$(2.8) \quad (1 - at)^{-\alpha/a} = (1 - at)^{-\beta/a} \sum_{m=0}^{\infty} \left(\frac{\alpha - \beta}{a}\right)_m \frac{(at)^m}{m!}$$

### 3. Generating Relations

We obtained some generating relations of (1.6) as

$$(3.1) \quad \sum_{n=0}^{\infty} T_{kn}^{(\alpha+s-1)}(x)t^n = (1 - t)^{-(\alpha+s)} \exp[p_k(x) - p_k\{x(1 - t)^{-1}\}]$$

$$\sum_{n=0}^{\infty} T_{kn}^{(\alpha-n+s-1)}(x)t^n = (1+t)^{-1+(\alpha+s)} \exp[p_k(x) - p_k\{x(1+t)\}]$$

(3.2)

$$\begin{aligned} & \sum_{m=0}^{\infty} \binom{m+n}{n} T_{k(n+m)}^{(\alpha+s-1)}(x) t^m \\ &= (1-t)^{-(\alpha+s+n)} \exp[p_k(x) - p_k\{x(1-t)^{-1}\}] T_{kn}^{(\alpha+s-1)}\{x(1-t)^{-1}\} \end{aligned}$$

(3.3)

$$\begin{aligned} & \sum_{m=0}^{\infty} \binom{m+n}{n} T_{k(n+m)}^{(\alpha-m+s-1)}(x) t^m \\ &= (1+t)^{\alpha+s-1} \exp[p_k(x) - p_k\{x(1+t)\}] T_{kn}^{(\alpha-m+s-1)}\{x(1+t)\} \end{aligned}$$

(3.4)

**Proof of (3.1).** From (1.6), we consider

$$\sum_{n=0}^{\infty} x^n T_{kn}^{(\alpha+s-1)}(x)t^n = x^{-\alpha} \exp\{p_k(x)\} e^{t\theta} [x^{\alpha} \exp\{-p_k(x)\}]$$

and using (2.6), above equation reduces to,

$$\begin{aligned} \sum_{n=0}^{\infty} x^n T_{kn}^{(\alpha-s+1)}(x)t^n &= x^{-\alpha} \exp\{p_k(x)\} x^{\alpha} (1-xt)^{-(\alpha+s)} \exp[-p_k\{x(1-xt)^{-1}\}] \\ &= (1-xt)^{-(\alpha+s)} \exp[p_k(x) - p_k\{x(1-xt)^{-1}\}] \end{aligned}$$

replacing  $t$  by  $t/x$ , which gives (3.1).

**Proof of (3.2).** From (1.6) we consider,

$$T_{kn}^{(\alpha-n+s-1)}(x) = \frac{1}{n!} x^{-(\alpha-n)-n} \exp\{p_k(x)\} \theta^n [x^{\alpha-n} \exp\{-p_k(x)\}]$$

or

$$\sum_{n=0}^{\infty} T_{kn}^{(\alpha-n+s-1)}(x) t^n = (x)^{-\alpha} \exp\{p_k(x)\} e^{t\theta} [x^{\alpha-n} \exp(-p_k(x))]$$

by using (2.7), we get

$$\begin{aligned} \sum_{n=0}^{\infty} T_{kn}^{(\alpha-n+s-1)}(x) t^n &= x^{-\alpha} \exp\{p_k(x)\} x^{\alpha} (1+t)^{-1+(\alpha+s)} \exp\{-p_k\{x(1+t)\}\} \\ &= (1+t)^{-1+(\alpha+s)} \exp[p_k(x) - p_k\{x(1+t)\}]. \end{aligned}$$

**Proof of (3.3).** Again from (1.6) we consider,

$$\theta^n [x^{\alpha} \exp\{-p_k(x)\}] = n! x^{\alpha+n} \exp\{-p_k(x)\} T_{kn}^{(\alpha+s-1)}(x)$$

or

$$e^{t\theta} (\theta^n [x^{\alpha} \exp\{-p_k(x)\}]) = n! e^{t\theta} [x^{\alpha+n} \exp\{-p_k(x)\} T_{kn}^{(\alpha+s-1)}(x)]$$

using (2.6) we get,

$$\begin{aligned} \sum_{m=0}^{\infty} \frac{t^m \theta^{m+n}}{m!} [x^{\alpha} \exp\{-p_k(x)\}] \\ = n! x^{\alpha+n} (1-xt)^{-(\alpha+s+n)} \exp[-p_k\{x(1-xt)^{-1}\}] T_{kn}^{(\alpha+s-1)}\{x(1-xt)^{-1}\} \end{aligned}$$

therefore, we get

$$\begin{aligned} & \sum_{m=0}^{\infty} \frac{1}{m! n!} (m+n)! x^{\alpha+m+n} \exp\{-p_k(x)\} T_{k(m+n)}^{(\alpha+s-1)}(x) t^m \\ &= x^{\alpha+n} (1-xt)^{-(\alpha+s+n)} \exp[-p_k\{x(1-xt)^{-1}\}] T_{kn}^{(\alpha+s-1)}\{x(1-xt)^{-1}\} \end{aligned}$$

hence above equation reduces to,

$$\begin{aligned} & \sum_{m=0}^{\infty} x^m \binom{m+n}{n} T_{k(m+n)}^{(\alpha+s-1)}(x) t^m \\ &= (1-xt)^{-(\alpha+s+n)} \exp[p_k(x) - p_k\{x(1-xt)^{-1}\}] T_{kn}^{(\alpha+s-1)}\{x(1-xt)^{-1}\} \end{aligned}$$

replacing  $t$  by  $t/x$ , which gives (3.3).

**Proof of (3.4).** Again from (1.6) we consider,

$$\theta^n [x^\alpha \exp\{-p_k(x)\}] = n! x^{\alpha+n} \exp\{-p_k(x)\} T_{kn}^{(\alpha+s-1)}(x)$$

replacing  $\alpha$  by  $\alpha - m$ , we get

$$\theta^n [x^{\alpha-m} \exp\{-p_k(x)\}] = n! x^{\alpha-m+n} \exp\{-p_k(x)\} T_{kn}^{(\alpha-m+s-1)}(x)$$

or

$$e^{t\theta} (\theta^n [x^{\alpha-m} E_\alpha\{-p_k(x)\}]) = n! e^{t\theta} [x^{(\alpha+n)-m} \exp\{-p_k(x)\} T_{kn}^{(\alpha-m+s-1)}(x)]$$

using (2.7) we get,

$$\begin{aligned} & \sum_{n=0}^{\infty} \frac{t^m \theta^{m+n}}{m!} [x^{\alpha-m} \exp\{-p_k(x)\}] \\ &= n! x^{\alpha+n} (1+t)^{\alpha+s-1} \exp[-p_k\{x(1+t)\}] T_{kn}^{(\alpha-m+s-1)}\{x(1+t)\} \end{aligned}$$

therefore, we get

$$\begin{aligned} \sum_{m=0}^{\infty} \frac{1}{m! n!} (m+n)! x^{\alpha-m+m+n} \exp\{-p_k(x)\} T_{k(m+n)}^{(\alpha-m+s-1)}(x) t^m \\ = x^{\alpha+n} (1+t)^{\alpha+s-1} \exp[-p_k\{x(1+t)\}] T_{kn}^{(\alpha-m+s-1)}\{x(1+t)\} \end{aligned}$$

which reduces to (3.4).

#### 4. Finite Summation Formulae

We obtained finite summation formula for (1.6) as

$$(4.1) \quad T_{kn}^{(\alpha+s-1)}(x) = \sum_{m=0}^n (m!)^{-1} (\alpha - \beta)_m T_{k(n-m)}^{(\beta+s-1)}(x)$$

$$(4.2) \quad T_{kn}^{(\alpha+s-1)}(x) = \sum_{m=0}^n \frac{1}{m!} (\alpha)_m T_{k(n-m)}^{(s-1)}(x)$$

**Proof of (4.1).** We can write (1.6) as,

$$\sum_{n=0}^{\infty} x^n T_{kn}^{(\alpha+s-1)}(x) t^n = x^{-\alpha} \exp\{p_k(x)\} e^{t\theta} [x^{\alpha} \exp\{-p_k(x)\}]$$

by using (2.6), we write

$$\begin{aligned} \sum_{n=0}^{\infty} x^n T_{kn}^{(\alpha+s-1)}(x) t^n \\ = x^{-\alpha} \exp\{p_k(x)\} x^{\alpha} (1 - xt)^{-(\alpha+s)} \exp[-p_k\{x(1 - xt)^{-1}\}] \\ = (1 - xt)^{-(\alpha+s)} \exp[p_k(x) - p_k\{x(1 - xt)^{-1}\}] \end{aligned}$$

applying (2.8), which yields

$$\sum_{n=0}^{\infty} x^n T_{kn}^{(\alpha+s-1)}(x) t^n$$



$$\begin{aligned}
 &= (1 - xt)^{-(\beta+s)} \sum_{m=0}^{\infty} (\alpha - \beta)_m \frac{(xt)^m}{m!} \exp[p_k(x) - p_k\{x(1 - xt)^{-1}\}] \\
 &= \sum_{n=0}^{\infty} (\alpha - \beta)_m \frac{x^m t^m}{m!} \exp\{p_k(x)\} (1 - xt)^{-(\beta+s)} \exp[-p_k\{x(1 - xt)^{-1}\}]
 \end{aligned}$$

using (3.1), above equation reduces to,

$$\begin{aligned}
 &\sum_{n=0}^{\infty} x^n T_{kn}^{(\alpha+s-1)}(x) t^n = \\
 &= \sum_{m=0}^{\infty} (\alpha - \beta)_m \frac{x^m t^m}{m!} \exp\{p_k(x)\} x^{-\beta} e^{t\theta} [x^{\beta} \exp\{-p_k(x)\}] \\
 &= \sum_{m,n=0}^{\infty} (\alpha - \beta)_m \frac{x^m t^{n+m}}{m! n!} \exp\{p_k(x)\} x^{-\beta} \theta^n [x^{\beta} \exp(-p_k(x))] \\
 &= \sum_{n=0}^{\infty} \sum_{m=0}^n \frac{1}{m!} (\alpha - \beta)_m \frac{x^{-\beta+m}}{(n - m)!} \exp\{p_k(x)\} \theta^{n-m} [x^{\beta} \exp\{-p_k(x)\}] t^n
 \end{aligned}$$

equating the coefficients of  $t^n$ , we get

$$x^n T_{kn}^{(\alpha+s-1)}(x) = \sum_{m=0}^n \frac{1}{m!} (\alpha - \beta)_m \frac{x^{-\beta+m}}{(n - m)!} \exp\{p_k(x)\} \theta^{n-m} [x^{\beta} \exp\{-p_k(x)\}]$$

Therefore, we obtain

$$T_{kn}^{(\alpha+s-1)}(x) = \sum_{m=0}^n \frac{1}{m!} (\alpha - \beta)_m \frac{x^{-\beta(-n-m)}}{(n - m)!} \exp\{p_k(x)\} \theta^{n-m} [x^{\beta} \exp\{-p_k(x)\}]$$

and applying (1.6) then above equation immediately leads to (4.1).

**Proof of (4.2).** We can write (1.6) as,

$$T_{kn}^{(\alpha+s-1)}(x) = \frac{1}{n!} x^{-\alpha-n} \exp\{p_k(x)\} \theta^n [x x^{\alpha-1} \exp\{-p_k(x)\}]$$

using (2.3) we get,

and by using (2.1) which yields,

$$\begin{aligned}
 T_{kn}^{(\alpha+s-1)}(x) &= \frac{1}{n!} x^{-\alpha-n} \exp\{p_k(x)\} x \sum_{m=0}^n \frac{n!}{m! (n-m)!} \\
 &\times x^{n-m} [(s+xD)(s+1+xD)(s+2+xD) \dots (s+(n-m-1)+xD)] \exp\{-p_k(x)\} \\
 &\times x^m [(1+xD)(2+xD)(3+xD) \dots (m+xD)] x^{\alpha-1} \\
 T_{kn}^{(\alpha+s-1)}(x) &= \exp\{p_k(x)\} \sum_{m=0}^n \frac{1}{m! (n-m)!} \prod_{i=0}^{n-m-1} (s+i+xD) \exp\{-p_k(x)\} (\alpha)_m \\
 (4.3)
 \end{aligned}$$

Putting  $\alpha = 0$  and replacing  $n$  by  $n - m$  in (1.6) which reduces to

$$T_{k(n-m)}^{(s-1)}(x) = \frac{1}{(n-m)!} x^{-(n-m)} \exp\{p_k(x)\} \theta^{n-m} [\exp\{-p_k(x)\}]$$

thus, we have

$$\frac{1}{(n-m)!} \theta^{n-m} [\exp\{-p_k(x)\}] = \frac{x^{n-m}}{\exp\{p_k(x)\}} T_{k(n-m)}^{(s-1)}(x)$$

using (2.1), we get

$$\begin{aligned}
 \frac{1}{(n-m)!} \prod_{i=0}^{n-m-1} (s+i+xD) [\exp\{-p_k(x)\}] &= \frac{1}{\exp\{p_k(x)\}} T_{k(n-m)}^{(s-1)}(x). \\
 (4.4)
 \end{aligned}$$

use of (4.4) and (4.3), gives complete proof of (4.2).

## 5. Concluding Remarks

Some special cases of  $T_{kn}^{(\alpha+s-1)}(x)$  polynomials are given below:

If we replace  $\alpha$  by  $\alpha + 1$ ,  $p_k(x) = p_1(x) = x$  and  $s = 0$  in (1.6), then this equation reduces to

$$(5.1) \quad T_n^{(\alpha)}(x) = L_n^{(\alpha)}(x) = Z_n^\alpha(x; 1) = Y_n^\alpha(x; 1).$$

Again replacing  $\alpha$  by  $\alpha + 1$ ,  $p_k(x) = px^r$  and  $s = 0$  in (1.6), which gives

$$(5.2) \quad T_{rn}^{(\alpha)}(x) = T_{rn}^{(\alpha)}(x, p).$$

Substituting  $\alpha = 1 - n$ ,  $p_k(x) = x^2$ ,  $s = 0$  in (1.6) and using (1.11), which yields

$$(5.3) \quad T_{2n}^{(1-n)}(x) = \frac{(-x)^n}{n!} H_n(x).$$

## Acknowledgements

We express our sincere thanks to the referees for their kind comments for the improvement of this manuscript.

## References

- [1] A. M. Chak, *A class of polynomials and generalization of stirling numbers*, Duke J. Math. **23** pp. 45–55, (1956).
- [2] S. K. Charttejea, *On generalization of Laguerre polynomials*, Rend. Mat. Univ. Padova **34**, pp. 180–190, (1964).
- [3] H. W. Gould and A. T. Hopper, *Operational formulas connected with two generalizations of Hermite polynomials*, Duke Math. J. **29**, pp. 51–63, (1962).
- [4] E. B. McBride, *Obtaining Generating Functions*, Springer-Verlag New York, (1971).
- [5] H. B. Mittal, *A generalization of Laguerre polynomials*, Publ. Math. Debrecen, **18**, pp. 53–88, (1971).

- [6] H. B. Mittal, *Operational representations for the generalized Laguerre polynomial*, Glasnik Mat. Ser III, **6**(26), pp. 45–53, (1971).
- [7] H. B. Mittal, *Bilinear and bilateral generating relations*, American J. Math. **99**, pp. 23–45, (1977).
- [8] K. R. Patil and N.K. Thakre, *Operational formulas for a function defined by a generalized Rodrigues formula-II*, Sci. J. Shivaji Univ. **15**, pp. 1–10, (1975).
- [9] E. D. Rainville, *Special Functions*, The Macmillan Company, New York, (1960).
- [10] R. P. Singh, *On generalized Truesdell polynomials*, Rivista de Matematica, **8**, pp. 345–353, (1968).
- [11] H. M. Srivastava, *Some biorthogonal polynomials suggested by the Laguerre polynomials*, Pacific J. Math. **98**(1), pp. 235–250, (1982).
- [12] H. M. Srivastava and H.L. Manocha, *A Treatise on Generating Functions*, Ellis Horwood Limited, Chichester, (1984).

**A. K. Shukla**

Department of Mathematics  
S. V. National Institute of Technology  
Surat-395007  
India  
e-mail : ajayshukla2@rediffmail.com

and

**J. C. Prajapati**

Department of Mathematics  
S. V. National Institute of Technology  
Surat-395007  
India  
e-mail : jyotindra18@rediffmail.com