

Proyecciones  
Vol. 22, N° 3, pp. 237-242, December 2003.  
Universidad Católica del Norte  
Antofagasta - Chile  
DOI: 10.4067/S0716-09172003000300005

# CLASSES OF FORMS WITT EQUIVALENT TO A SECOND TRACE FORM OVER FIELDS OF CHARACTERISTIC TWO\*

*ANA CECILIA DE LA MAZA*  
*Universidad de Talca, Chile*

## Abstract

*Let  $F$  be a field of characteristic two. We determine all non-hyperbolic quadratic forms over  $F$  that are Witt equivalent to a second trace form.*

**Mathematics Subject Classification :** *Primary 11 E 04; Secondary 11 E 81*

**Key Words:** *Trace form, characteristic two*

---

\*This work was supported by Fondecyt grants N°1010205, and by Programa formas extremas y representación de formas cuadráticas, Universidad de Talca

## 1. Introduction

Let  $E/F$  be a finite separable field extension. We define the trace form for this extension by  $q(x) = \text{tr}_{E/F}(x^2)$ . When the characteristic of  $F$  is not equal to 2, the trace form  $(E, q)$  is non-degenerate. However, if the characteristic is 2 then  $(E, q)$  is degenerate and splits as  $[1] \perp V$ , with  $V$  totally isotropic. It is therefore natural to introduce a modified “second trace form”. To this end one considers for each  $a \in E$  its characteristic polynomial

$$(1.1) \quad p(x, a) = x^n - T_1(a)x^{n-1} + T_2(a)x^{n-2} + \cdots + (-1)^n T_n(a)$$

(whence  $T_1(a) = \text{tr}_{E/F}(a)$  and  $T_n(a)$  is the norm of  $a$ ). It is clear that  $(E, T_2)$  is a quadratic form. When the degree  $n$  of the extension is odd this form is necessarily singular. To arrive at a non-degenerate form, two methods have been proposed in the literature. One method, due to Bergé and Martinet [BM], increases the dimension of the space by 1 using the étale  $F$ -algebra. The other method, due to Revoy [R], reduces the dimension of the space by 1. In this note we will adopt the second method and call such forms *2-trace forms*.

We consider the following problem: Which elements  $[q] \neq 0$  of the Witt-group  $W_q(F)$  are represented by 2-trace forms?. Our theorem 3 fully answers this question. Moreover, we will partially answer the same question for  $[q] = 0$  (see Prop. 1 and Prop. 2). For fields of characteristic not equal to 2 this problem seems quite more complicated: for partial results concerning generic fields one may consult [CP] and [EHP]; a complete solution for Hilbertian fields is given in [Sch], [KS] and [Wat].

## 2. The second trace form

As we remarked in the introduction, there are two ways to define a second trace form. In this section we will prove that in fact the corresponding forms are Witt equivalent.

Let  $E/F$  be a finite separable field extension. The second trace form  $T_{E/F}$  of the extension  $E/F$  was defined by Revoy [R] as  $(E, T_2)$  if the degree  $[E : F]$  is even, and as  $(E_0, T_2)$  if the degree is odd, where  $T_1, T_2$  are given by (1.1) and  $E_0 = \text{Ker } T_1$ . It is important to remark that the bilinear form  $b_q$  associated to  $T_{E/F}$  satisfies the following relations:

$$(2.1) \quad b_q(x, y) = T_2(x + y) - T_2(x) - T_2(y) = T_1(xy) - T_1(x)T_1(y),$$

$$(2.2) \quad T_1(x^2) = (T_1(x))^2 \quad \text{and} \quad b_q(x^2, y^2) = b_q(x, y)^2$$

On the other hand, Bergé and Martinet defined in [BM] the second trace form as Revoy if the degree  $[E : F]$  is even and as  $(E \times F, T_2)$  if not.

**Theorem 1.** *The Revoy form and the Bergé-Martinet form are Witt equivalent.*

**Proof.** If  $[E : F]$  is odd, then the form  $(E \times F, T_2)$  of Bergé and Martinet [BM, p. 14] splits as follows  $(F(1, 0) + F(0, 1)) \perp E_0 \times \{0\}$ . Since  $F(1, 0) + F(0, 1)$  is an hyperbolic plane, the claim follows immediately.  $\square$

### 3. 2-algebraic forms

In this section we determine all non-hyperbolic quadratic forms over  $F$  that are Witt equivalent to some second trace form. Furthermore we give fields where hyperbolic forms are Witt equivalent to a second trace form.

**Theorem 2.** *Let  $E/F$  be a finite separable field extension, with  $[E : F] = 2n + 1$  or  $[E : F] = 2n$ . Then  $T_{E/F} = (n - 1)\mathbf{H} \perp [1, a]$ , for some  $a \in F$ .*

**Proof.** The assertion is deduced from Theorem 1 above and Theorem 3.5 in [BM, p. 13-14]. In order to illustrate the ideas, we will give the proof in case  $[E : F] = 2n + 1$ .

Since  $(E_0, T_2)$  is nonsingular, there exists a symplectic basis  $\{e_i, f_i\}_{1 \leq i \leq n}$  of  $E_0$ . Hence for each  $1 \leq i \leq n$  there exists  $x_i, y_i \in Fe_i + Ff_i$  such that  $T_2(x_i) \neq 0$  and  $b_q(x_i, y_i) = 1$  (note that  $[0, 0] = [1, 0]$  [Sa1, p. 150]).

Put  $e'_i := T_2(x_i)^{-1}x_i^2$  and  $f'_i := T_2(x_i)y_i^2$ . We have by (2.2) that  $e'_i, f'_i \in E_0$  and  $b_q(e'_i, f'_i) = 1$ . Since  $T_2(e'_i) = 1$ , we obtain a symplectic basis such that the quadratic form decomposes as follows

$$T_{E/F} = [1, a_1] + [1, a_2] + \cdots + [1, a_n],$$

where  $a_i = T_2(f'_i)$ . Using the relation  $[1, b] + [1, c] \cong [1, b + c] + \mathbf{H}$  over  $F$ , we obtain  $T_{E/F} \cong (n - 1)\mathbf{H} + [1, a]$  with  $a \equiv a_1 + a_2 + \cdots + a_n \pmod{\wp(F)}$ , where  $\wp(F) := \{x^2 + x \mid x \in F\}$ .  $\square$

A form  $T$  over  $F$  is called 2-algebraic if it is Witt equivalent to some second trace form.

**Corollary 1.** *A non hyperbolic quadratic form  $(V, q)$  over  $F$  is 2-algebraic if and only if  $(V, q) = r\mathbf{H} \perp (V_a, q_a)$ , with  $V_a$  an anisotropic plane representing 1.*

**Proof:**  $\Rightarrow$ ) Is clear by Theorem 2.

$\Leftarrow$ ) Let  $(V, q) = r\mathbf{H} \perp (V_a, q_a)$ , with  $(V_a, q_a)$  2-dimensional anisotropic representing 1. Then we can rewrite  $(V_a, q_a) = [1, b]$ , where  $b \notin \wp(F)$ . Let  $E = F(\alpha)$  with  $\alpha \in \overline{F}$  and  $\alpha^2 + \alpha + b = 0$ . We have that  $(E, T_{E/F}) = [1, b]$ .  $\square$

**Example 1.** Let  $F = \mathbf{F}_2(a)$  and  $E = F(b)$ , where  $a^2 + a + 1 = 0$  and  $b^3 + b + a = 0$ . Then  $T_{E/F} = [1, a] \neq [1, 1]$ .

In fact, using (2.1), we see that  $\{b^2, (1+a)b\}$  is a symplectic basis for  $E_0 = \text{Ker } T_1$ . Since  $p(x, (1+a)b) = x^3 + ax + a$ ,  $1 \in \wp(F)$  and  $a \notin \wp(F)$ , we obtain the form  $(E_0, T_2) = [1, a] \neq [1, 1]$ .

**Corollary 2.** *If a non hyperbolic quadratic form  $(V, q)$  over  $F$  is 2-algebraic then there exists a quadratic extension field  $E$  of  $F$  such that the extension  $(V \otimes_F E, q_E)$  is hyperbolic.*

**Proof:** See the proof of Theorem 3 and note that  $[1, b] = \mathbf{H}$  over  $E = F(\alpha)$ , with  $\alpha^2 + \alpha + b = 0$  (see [Sa1, p. 150]).  $\square$

**Theorem 3.** *Let  $F = \mathbf{F}_2$  or  $F = \mathbf{F}_2(t)$  with  $t$  transcendental over  $\mathbf{F}_2$ . Then the hyperbolic quadratic forms over  $F$  are 2-algebraic.*

**Proof.** We only need to find an extension  $E$  of  $F$  such that  $T_{E/F}$  is hyperbolic. We first remark that  $p(x) := x^4 + x^3 + 1$  is irreducible over  $\mathbf{F}_2$  and also over  $\mathbf{F}_2(t)$ . Let  $\alpha$  be a root of  $p$  and  $E = F(\alpha)$ . We decompose the trace form  $T_{E/F}$  with respect to the basis  $\{\alpha, 1 + \alpha^3\} \cup \{\alpha^2, \alpha + \alpha^2 + \alpha^3\}$ . Noting that this basis has the elements conjugate to  $\alpha$ , it is easy to recognise that each vector basis is isotropic, and furthermore by (2.1) we see that it is a symplectic basis. Hence, the space is hyperbolic.  $\square$

**Theorem 4.** *Let  $F$  be a field. If there exists  $a \in F^*$  and  $n$  odd such that the polynomial  $x^n - a$  is irreducible over  $F[x]$ , then the hyperbolic quadratics space over  $F$  are 2-algebraic.*

**Proof.** Let  $E = F(\alpha)$ , where  $\alpha \in \overline{F}$  and  $\alpha^n = a$ . For  $1 \leq k \leq n-1$ , the linear transformation  $f_{\alpha^k} : x \mapsto x\alpha^k$  is given by the matrix  $c_{ij}(k)$ , where

$$c_{ij}(k) = \begin{cases} 1 & \text{if } j = i - k \\ a & \text{if } j = n + i - k \\ 0 & \text{otherwise} \end{cases}$$

Then for  $1 \leq k \leq n-1$ ,  $\alpha^k \in E_0$ , because  $c_{ii}(k) = 0$  for each  $i$ . Noting that  $T_1(a) = a$  we obtain the decomposition

$$E_0 = \langle \alpha, \alpha^{n-1} \rangle \perp \langle \alpha^2, \alpha^{n-2} \rangle \perp \cdots \perp \langle \alpha^{\frac{n-1}{2}}, \alpha^{\frac{n+1}{2}} \rangle,$$

where  $\langle x, y \rangle$  is the space generate by  $x$  and  $y$ . Hence, using that  $n \neq 2k$ , we deduce that  $T_2(\alpha^k) = 0$  for  $1 \leq k \leq n-1$ , so  $(E_0, T_2) = (\frac{n-1}{2})\mathbf{H}$ .  $\square$

## References

- [BM] Bergé, A.M., Martinet, J.: *Formes quadratiques et extensions en caractéristique 2*, Sémin. Théorie des Nombres de Bordeaux, exposé 35, (1982)-(1983).
- [CP] Conner, P., Perlis, R.: *A survey of trace forms of algebraic number fields*, World Scientific, Singapur, (1984).
- [EHP] Estes, R., Hurrelbrink, J., Perlis, R.: *Total positivity and algebraic Witt classes*, Comment. Math. Helvetici 60, pp. 284-290, (1985).
- [KS] Krüspkeemper, M., Scharlau, W.: *On trace forms of Hilbertian fields*, Comment. Math. Helvetici 63, pp. 296-304, (1988).
- [R] Revoy, P.: *Remarques sur la forme trace*, Linear and multilinear algebra 10, pp. 223-233, (1981).

- [Sa1] Sah, C.H.: *Symmetric bilinear forms and quadratic forms*, Journal of algebra 20, pp. 144-169, (1972).
- [Se] Serre, J.P.: *Linvariant de Witt de la forme  $\text{tr}(x^2)$* , Comment. Math. Helvetici 59, pp. 657-676, (1984).
- [Sch] Scharlau, W.: *On trace forms of algebraic number fields*, Math. Zeit. 127, pp. 125-127, (1987).
- [Wat] Waterhouse, W.: *Scaled trace forms over number fields*, Arch. Math. 47, pp. 229-231, (1986).

Received : January, 2003.

**Ana Cecilia de la Maza**

Instituto de Matemática y Física

Universidad de Talca

Casilla 747

Talca

e-mail : anace@inst-mat.otalca.cl