

Proyecciones Journal of Mathematics
Vol. 37, N° 1, pp. 133-151, March 2018.
Universidad Católica del Norte
Antofagasta - Chile

Some new Ostrowski type fractional integral inequalities for generalized (s, m, φ) -preinvex functions via Caputo k -fractional derivatives

Artion Kashuri

University "Ismail Qemali", Albania

and

Rozana Liko

University "Ismail Qemali", Albania

Received : June 2017. Accepted : July 2017

Abstract

In the present paper, the notion of generalized (s, m, φ) -preinvex function is applied to establish some new generalizations of Ostrowski type integral inequalities via Caputo k -fractional derivatives. At the end, some applications to special means are given.

2010 Mathematics Subject Classification : Primary: 26A51;
Secondary: 26A33, 26D07, 26D10, 26D15.

Keywords : *Ostrowski type inequality, Hölder's inequality, power mean inequality, s -convex function in the second sense, m -invex.*

1. Introduction and Preliminaries

The following notations are used throughout this paper. We use I to denote an interval on the real line $\mathbf{R} = (-\infty, +\infty)$ and I° to denote the interior of I . For any subset $K \subseteq \mathbf{R}^n$, K° is used to denote the interior of K . \mathbf{R}^n is used to denote a n -dimensional vector space. The set of continuous differentiable functions of order n on the interval $[a, b]$ is denoted by $C^n[a, b]$.

The following result is known in the literature as the Ostrowski inequality (see [10]), which gives an upper bound for the approximation of the integral average $\frac{1}{b-a} \int_a^b f(t)dt$ by the value $f(x)$ at point $x \in [a, b]$.

Theorem 1.1. *Let $f : I \rightarrow \mathbf{R}$, where $I \subseteq \mathbf{R}$ is an interval, be a mapping differentiable in the interior I° of I , and let $a, b \in I^\circ$ with $a < b$. If $|f'(x)| \leq M$ for all $x \in [a, b]$, then*

$$\left| f(x) - \frac{1}{b-a} \int_a^b f(t)dt \right| \leq M(b-a) \left[\frac{1}{4} + \frac{\left(x - \frac{a+b}{2} \right)^2}{(b-a)^2} \right], \quad \forall x \in [a, b]. \quad (1.1)$$

For other recent results concerning Ostrowski type inequalities (see [1]-[3],[10],[16],[34]-[38],[42]). Ostrowski inequality is playing a very important role in all the fields of mathematics, especially in the theory of approximations. Thus such inequalities were studied extensively by many researches and numerous generalizations, extensions and variants of them for various kind of functions like bounded variation, synchronous, Lipschitzian, monotonic, absolutely, continuous and n -times differentiable mappings etc. appeared in a number of papers (see [4]-[15],[17]-[20]). In recent years, one more dimension has been added to this studies, by introducing a number of integral inequalities involving various fractional operators like Riemann-Liouville, Erdelyi-Kober, Katugampola, conformable fractional integral operators etc. by many authors (see [21]-[31]). Riemann-Liouville fractional integral operators are the most central between these fractional operators.

In numerical analysis many quadrature rules have been established to approximate the definite integrals. Ostrowski inequality provides the bounds of many numerical quadrature rules (see [8]). In recent decades Ostrowski inequality is studied in fractional calculus point of view by many mathematicians (see [9]-[11]).

Definition 1.2. (see [33]) A function $f : [0, +\infty) \rightarrow \mathbf{R}$ is said to be s -convex in the second sense, if

$$(1.2) \quad f(\lambda x + (1 - \lambda)y) \leq \lambda^s f(x) + (1 - \lambda)^s f(y)$$

for all $x, y \geq 0$, $\lambda \in [0, 1]$ and $s \in (0, 1]$.

It is clear that a 1-convex function must be convex on $[0, +\infty)$ as usual. The s -convex functions in the second sense have been investigated in (see [33]).

Definition 1.3. (see [39]) A set $K \subseteq \mathbf{R}^n$ is said to be invex with respect to the mapping $\eta : K \times K \rightarrow \mathbf{R}^n$, if $x + t\eta(y, x) \in K$ for every $x, y \in K$ and $t \in [0, 1]$.

Notice that every convex set is invex with respect to the mapping $\eta(y, x) = y - x$, but the converse is not necessarily true. For more details (see [39],[40]).

Definition 1.4. (see [41]) The function f defined on the invex set $K \subseteq \mathbf{R}^n$ is said to be preinvex with respect η , if for every $x, y \in K$ and $t \in [0, 1]$, we have that

$$f(x + t\eta(y, x)) \leq (1 - t)f(x) + tf(y).$$

The concept of preinvexity is more general than convexity since every convex function is preinvex with respect to the mapping $\eta(y, x) = y - x$, but the converse is not true.

Definition 1.5. For $k \in \mathbf{R}^+$ and $x \in \mathbf{C}$, the k -gamma function is defined by

$$(1.3) \quad \Gamma_k(x) = \lim_{n \rightarrow \infty} \frac{n! k^n n k^{\frac{x}{k}-1}}{(x)_{n,k}}.$$

Its integral representation is given by

$$(1.4) \quad \Gamma_k(\alpha) = \int_0^\infty t^{\alpha-1} e^{-\frac{t^k}{k}} dt.$$

One can note that $\Gamma_k(\alpha + k) = \alpha\Gamma_k(\alpha)$.

For $k = 1$, (1.4) gives integral representation of gamma function.

Definition 1.6. For $k \in \mathbf{R}^+$ and $x \in \mathbf{C}$, the k -beta function with two parameters x and y is defined as

$$(1.5) \quad \beta_k(x, y) = \frac{1}{k} \int_0^1 t^{\frac{x}{k}-1} (1-t)^{\frac{y}{k}-1} dt.$$

For $k = 1$, (1.5) gives integral representation of beta function.

Theorem 1.7. Let $x, y > 0$, then for k -gamma and k -beta function the following equality holds

$$(1.6) \quad \beta_k(x, y) = \frac{\Gamma_k(x)\Gamma_k(y)}{\Gamma_k(x+y)}.$$

Definition 1.8. (see [12]) Let $\alpha > 0$ and $\alpha \notin \{1, 2, 3, \dots\}$, $n = [\alpha] + 1$, $f \in C^n[a, b]$ such that $f^{(n)}$ exists and are continuous on $[a, b]$. The Caputo fractional derivatives of order α are defined as follows:

$$(1.7) \quad {}^cD_{a+}^\alpha f(x) = \frac{1}{\Gamma(n-\alpha)} \int_a^x \frac{f^{(n)}(t)}{(x-t)^{\alpha-n+1}} dt, \quad x > a$$

and

$$(1.8) \quad {}^cD_{b-}^\alpha f(x) = \frac{(-1)^n}{\Gamma(n-\alpha)} \int_x^b \frac{f^{(n)}(t)}{(t-x)^{\alpha-n+1}} dt, \quad x < b.$$

If $\alpha = n \in \{1, 2, 3, \dots\}$ and usual derivative of order n exists, then Caputo fractional derivative $({}^cD_{a+}^\alpha f)(x)$ coincides with $f^{(n)}(x)$. In particular we have

$$(1.9) \quad \left({}^cD_{a+}^0 f\right)(x) = \left({}^cD_{b-}^0 f\right)(x) = f(x)$$

where $n = 1$ and $\alpha = 0$.

In the following we recall Caputo k -fractional derivatives.

Definition 1.9. (see [13]) Let $\alpha > 0$, $k \geq 1$ and $\alpha \notin \{1, 2, 3, \dots\}$, $n = [\alpha] + 1$, $f \in C^n[a, b]$. The Caputo k -fractional derivatives of order α are defined as follows:

$$(1.10) \quad {}^cD_{a+}^{\alpha,k} f(x) = \frac{1}{k\Gamma_k(n-\frac{\alpha}{k})} \int_a^x \frac{f^{(n)}(t)}{(x-t)^{\frac{\alpha}{k}-n+1}} dt, \quad x > a$$

and

$$(1.11) \quad {}^cD_{b-}^{\alpha,k} f(x) = \frac{(-1)^n}{k\Gamma_k(n-\frac{\alpha}{k})} \int_x^b \frac{f^{(n)}(t)}{(t-x)^{\frac{\alpha}{k}-n+1}} dt, \quad x < b.$$

The aim of this paper is to establish some generalizations of Ostrowski type inequalities using new identity given in Section 2 to generalized (s, m, φ) -preinvex functions via Caputo k -fractional derivatives. In Section 3, some applications to special means are given. In Section 4, some conclusions and future research are also given.

2. Main results

Definition 2.1. (see [32]) A set $K \subseteq \mathbf{R}^n$ is said to be m -invex with respect to the mapping $\eta : K \times K \times (0, 1] \rightarrow \mathbf{R}^n$ for some fixed $m \in (0, 1]$, if $mx + t\eta(y, x, m) \in K$ holds for each $x, y \in K$ and any $t \in [0, 1]$.

Remark 2.2. In Definition 2.1, under certain conditions, the mapping $\eta(y, x, m)$ could reduce to $\eta(y, x)$. For example when $m = 1$, then the m -invex set degenerates an invex set on K .

Definition 2.3. (see [2]) Let $K \subseteq \mathbf{R}$ be an open m -invex set with respect to $\eta : K \times K \times (0, 1] \rightarrow \mathbf{R}$ and $\varphi : I \rightarrow K$ a continuous function. For $f : K \rightarrow \mathbf{R}$ and any fixed $s, m \in (0, 1]$, if

$$(2.1) \quad f(m\varphi(x) + t\eta(\varphi(y), \varphi(x), m)) \leq m(1-t)^s f(\varphi(x)) + t^s f(\varphi(y))$$

is valid for all $x, y \in I, t \in [0, 1]$, then we say that $f(x)$ is a generalized (s, m, φ) -preinvex function with respect to η .

Remark 2.4. In Definition 2.3, it is worthwhile to note that the class of generalized (s, m, φ) -preinvex function is a generalization of the class of s -convex in the second sense function given in Definition 1.2.

In this section, in order to present some new Ostrowski type integral inequalities to generalized (s, m, φ) -preinvex functions via Caputo k -fractional derivatives, we need the following new interesting lemma to obtain our results.

Lemma 2.5. Let $\alpha > 0, k \geq 1$ and $\alpha \notin \{1, 2, 3, \dots\}$, $n = [\alpha] + 1$. Let $\varphi : I \rightarrow K$ be a continuous function. Suppose $K \subseteq \mathbf{R}$ be an open

m-invex subset with respect to $\eta : K \times K \times (0, 1] \rightarrow \mathbf{R}$ for any fixed $m \in (0, 1]$. Assume that $f : K \rightarrow \mathbf{R}$ is a function on K° such that $f \in C^{n+1}[m\varphi(a), m\varphi(a) + \eta(\varphi(b), \varphi(a), m)]$, where $\eta(\varphi(b), \varphi(a), m) > 0$. Then we have the following equality for Caputo k-fractional derivatives

$$\begin{aligned}
& \frac{\eta^{n-\frac{\alpha}{k}}(\varphi(x), \varphi(a), m) f^{(n)}(m\varphi(a) + \eta(\varphi(x), \varphi(a), m))}{\eta(\varphi(b), \varphi(a), m)} \\
& - \frac{\eta^{n-\frac{\alpha}{k}}(\varphi(x), \varphi(b), m) f^{(n)}(m\varphi(b) + \eta(\varphi(x), \varphi(b), m))}{\eta(\varphi(b), \varphi(a), m)} \\
& + (-1)^{n+1} \frac{(nk - \alpha)\Gamma_k(n - \frac{\alpha}{k})}{\eta(\varphi(b), \varphi(a), m)} \\
& \times \left[{}^c D_{(m\varphi(a) + \eta(\varphi(x), \varphi(a), m))^-}^{\alpha, k} f(m\varphi(a)) - {}^c D_{(m\varphi(b) + \eta(\varphi(x), \varphi(b), m))^-}^{\alpha, k} f(m\varphi(b)) \right] \\
& = \frac{\eta^{n-\frac{\alpha}{k}+1}(\varphi(x), \varphi(a), m)}{\eta(\varphi(b), \varphi(a), m)} \int_0^1 t^{n-\frac{\alpha}{k}} f^{(n+1)}(m\varphi(a) + t\eta(\varphi(x), \varphi(a), m)) dt \\
& - \frac{\eta^{n-\frac{\alpha}{k}+1}(\varphi(x), \varphi(b), m)}{\eta(\varphi(b), \varphi(a), m)} \int_0^1 t^{n-\frac{\alpha}{k}} f^{(n+1)}(m\varphi(b) + t\eta(\varphi(x), \varphi(b), m)) dt.
\end{aligned}$$

Proof. Throughout this paper we denote

$$\begin{aligned}
& I_{f, \eta, \varphi}(x; \alpha, k, n, m, a, b) \\
& = \frac{\eta^{n-\frac{\alpha}{k}+1}(\varphi(x), \varphi(a), m)}{\eta(\varphi(b), \varphi(a), m)} \int_0^1 t^{n-\frac{\alpha}{k}} f^{(n+1)}(m\varphi(a) + t\eta(\varphi(x), \varphi(a), m)) dt \\
& - \frac{\eta^{n-\frac{\alpha}{k}+1}(\varphi(x), \varphi(b), m)}{\eta(\varphi(b), \varphi(a), m)} \int_0^1 t^{n-\frac{\alpha}{k}} f^{(n+1)}(m\varphi(b) + t\eta(\varphi(x), \varphi(b), m)) dt.
\end{aligned} \tag{2.2}$$

Integrating by parts, we get

$$\begin{aligned}
I_{f, \eta, \varphi}(x; \alpha, k, n, m, a, b) & = \frac{\eta^{n-\frac{\alpha}{k}+1}(\varphi(x), \varphi(a), m)}{\eta(\varphi(b), \varphi(a), m)} \\
& \times \left[\frac{t^{n-\frac{\alpha}{k}} f^{(n)}(m\varphi(a) + t\eta(\varphi(x), \varphi(a), m))}{\eta(\varphi(x), \varphi(a), m)} \right]_0^1
\end{aligned}$$

$$\begin{aligned}
 & -\frac{n-\frac{\alpha}{k}}{\eta(\varphi(x), \varphi(a), m)} \int_0^1 t^{n-\frac{\alpha}{k}-1} f^{(n)}(m\varphi(a) + t\eta(\varphi(x), \varphi(a), m)) dt \\
 & \quad - \frac{\eta^{n-\frac{\alpha}{k}+1}(\varphi(x), \varphi(b), m)}{\eta(\varphi(b), \varphi(a), m)} \\
 & \quad \times \left[\frac{t^{n-\frac{\alpha}{k}} f^{(n)}(m\varphi(b) + t\eta(\varphi(x), \varphi(b), m))}{\eta(\varphi(x), \varphi(b), m)} \right]_0^1 \\
 & -\frac{n-\frac{\alpha}{k}}{\eta(\varphi(x), \varphi(b), m)} \int_0^1 t^{n-\frac{\alpha}{k}-1} f^{(n)}(m\varphi(b) + t\eta(\varphi(x), \varphi(b), m)) dt \\
 & = \frac{\eta^{n-\frac{\alpha}{k}}(\varphi(x), \varphi(a), m) f^{(n)}(m\varphi(a) + \eta(\varphi(x), \varphi(a), m))}{\eta(\varphi(b), \varphi(a), m)} \\
 & \quad - \frac{\eta^{n-\frac{\alpha}{k}}(\varphi(x), \varphi(b), m) f^{(n)}(m\varphi(b) + \eta(\varphi(x), \varphi(b), m))}{\eta(\varphi(b), \varphi(a), m)} \\
 & \quad + (-1)^{n+1} \frac{(nk-\alpha)\Gamma_k(n-\frac{\alpha}{k})}{\eta(\varphi(b), \varphi(a), m)} \\
 & \times \left[{}^c D_{(m\varphi(a)+\eta(\varphi(x), \varphi(a), m))-}^{\alpha, k} f(m\varphi(a)) - {}^c D_{(m\varphi(b)+\eta(\varphi(x), \varphi(b), m))-}^{\alpha, k} f(m\varphi(b)) \right]. \\
 & \square
 \end{aligned}$$

By using Lemma 2.5, one can extend to the following results.

Theorem 2.6. Let $\alpha > 0$, $k \geq 1$ and $\alpha \notin \{1, 2, 3, \dots\}$, $n = [\alpha] + 1$. Let $\varphi : I \rightarrow K$ be a continuous function. Suppose $K \subseteq \mathbf{R}$ be an open m -invex subset with respect to $\eta : K \times K \times (0, 1] \rightarrow \mathbf{R}$ for any fixed $s, m \in (0, 1]$. Assume that $f : K \rightarrow \mathbf{R}$ is a function on K° such that $f \in C^{n+1}[m\varphi(a), m\varphi(a) + \eta(\varphi(b), \varphi(a), m)]$, where $\eta(\varphi(b), \varphi(a), m) > 0$. If $|f^{(n+1)}|$ is generalized (s, m, φ) -preinvex function on $[m\varphi(a), m\varphi(a) + \eta(\varphi(b), \varphi(a), m)]$ and $|f^{(n+1)}| \leq M$, then the following inequality for Caputo k -fractional derivatives holds:

$$\begin{aligned}
 |I_{f, \eta, \varphi}(x; \alpha, k, n, m, a, b)| & \leq M \left(m\beta \left(n - \frac{\alpha}{k} + 1, s + 1 \right) + \frac{1}{n + s - \frac{\alpha}{k} + 1} \right) \\
 (2.3) \quad & \times \left[\frac{|\eta(\varphi(x), \varphi(a), m)|^{n-\frac{\alpha}{k}+1} + |\eta(\varphi(x), \varphi(b), m)|^{n-\frac{\alpha}{k}+1}}{\eta(\varphi(b), \varphi(a), m)} \right].
 \end{aligned}$$

Proof. From Lemma 2.5, generalized (s, m, φ) -preinvexity of $|f^{(n+1)}|$, upper bound of $|f^{(n+1)}|$ and properties of the modulus, we have

$$\begin{aligned}
& |I_{f,\eta,\varphi}(x; \alpha, k, n, m, a, b)| \\
& \leq \frac{|\eta(\varphi(x), \varphi(a), m)|^{n-\frac{\alpha}{k}+1}}{\eta(\varphi(b), \varphi(a), m)} \int_0^1 t^{n-\frac{\alpha}{k}} |f^{(n+1)}(m\varphi(a) + t\eta(\varphi(x), \varphi(a), m))| dt \\
& \quad + \frac{|\eta(\varphi(x), \varphi(b), m)|^{n-\frac{\alpha}{k}+1}}{\eta(\varphi(b), \varphi(a), m)} \int_0^1 t^{n-\frac{\alpha}{k}} |f^{(n+1)}(m\varphi(b) + t\eta(\varphi(x), \varphi(b), m))| dt \\
& \leq \frac{|\eta(\varphi(x), \varphi(a), m)|^{n-\frac{\alpha}{k}+1}}{\eta(\varphi(b), \varphi(a), m)} \int_0^1 t^{n-\frac{\alpha}{k}} \left[m(1-t)^s |f^{(n+1)}(\varphi(a))| + t^s |f^{(n+1)}(\varphi(x))| \right] dt \\
& \quad + \frac{|\eta(\varphi(x), \varphi(b), m)|^{n-\frac{\alpha}{k}+1}}{\eta(\varphi(b), \varphi(a), m)} \int_0^1 t^{n-\frac{\alpha}{k}} \left[m(1-t)^s |f^{(n+1)}(\varphi(b))| + t^s |f^{(n+1)}(\varphi(x))| \right] dt \\
& \leq M \left(m\beta \left(n - \frac{\alpha}{k} + 1, s + 1 \right) + \frac{1}{n + s - \frac{\alpha}{k} + 1} \right) \\
& \quad \times \left[\frac{|\eta(\varphi(x), \varphi(a), m)|^{n-\frac{\alpha}{k}+1} + |\eta(\varphi(x), \varphi(b), m)|^{n-\frac{\alpha}{k}+1}}{\eta(\varphi(b), \varphi(a), m)} \right].
\end{aligned}$$

□

Corollary 2.7. Under the same conditions as in Theorem 2.6, if we choose $m = k = 1$, $\eta(\varphi(y), \varphi(x), m) = \varphi(y) - m\varphi(x)$ and $\varphi(x) = x$, $\forall x \in I$, we get the following inequality for Caputo fractional derivatives:

$$\begin{aligned}
& \left| \left[\frac{(x-a)^{n-\alpha} - (x-b)^{n-\alpha}}{b-a} \right] f^{(n)}(x) \right. \\
& \quad \left. + (-1)^{n+1} \frac{\Gamma(n-\alpha+1)}{b-a} \left[{}^c D_{x-}^\alpha f(a) - {}^c D_{x-}^\alpha f(b) \right] \right| \\
& \leq M \left(\beta(n-\alpha+1, s+1) + \frac{1}{n+s-\alpha+1} \right) \left[\frac{(x-a)^{n-\alpha+1} + (b-x)^{n-\alpha+1}}{b-a} \right]. \tag{2.4}
\end{aligned}$$

Theorem 2.8. Let $\alpha > 0$, $k \geq 1$ and $\alpha \notin \{1, 2, 3, \dots\}$, $n = [\alpha] + 1$. Let $\varphi : I \rightarrow K$ be a continuous function. Suppose $K \subseteq \mathbf{R}$ be an open m -invex subset with respect to $\eta : K \times K \times (0, 1) \rightarrow \mathbf{R}$ for any fixed $s, m \in (0, 1]$. Assume that $f : K \rightarrow \mathbf{R}$ is a function on K° such that $f \in C^{n+1}[m\varphi(a), m\varphi(a) + \eta(\varphi(b), \varphi(a), m)]$, where $+\eta(\varphi(b), \varphi(a), m) > 0$. If $|f^{(n+1)}|^q$ is generalized (s, m, φ) -preinvex function on $[m\varphi(a), m\varphi(a) + \eta(\varphi(b), \varphi(a), m)]$ $q > 1$, $p^{-1} + q^{-1} = 1$ and $|f^{(n+1)}| \leq M$, then the following inequality for Caputo k -fractional derivatives holds:

$$(2.5) \quad |I_{f,\eta,\varphi}(x; \alpha, k, n, m, a, b)| \leq \frac{M}{\left((n - \frac{\alpha}{k}) p + 1 \right)^{\frac{1}{p}}} \left(\frac{m+1}{s+1} \right)^{\frac{1}{q}} \times \left[\frac{|\eta(\varphi(x), \varphi(a), m)|^{n-\frac{\alpha}{k}+1} + |\eta(\varphi(x), \varphi(b), m)|^{n-\frac{\alpha}{k}+1}}{\eta(\varphi(b), \varphi(a), m)} \right].$$

Proof. Suppose that $q > 1$. From Lemma 2.5, generalized (s, m, φ) -preinvexity of $|f^{(n+1)}|^q$, Hölder inequality, upper bound of $|f^{(n+1)}|$ and properties of the modulus, we have

$$\begin{aligned} & |I_{f,\eta,\varphi}(x; \alpha, k, n, m, a, b)| \\ & \leq \frac{|\eta(\varphi(x), \varphi(a), m)|^{n-\frac{\alpha}{k}+1}}{\eta(\varphi(b), \varphi(a), m)} \int_0^1 t^{n-\frac{\alpha}{k}} |f^{(n+1)}(m\varphi(a) + t\eta(\varphi(x), \varphi(a), m))| dt \\ & + \frac{|\eta(\varphi(x), \varphi(b), m)|^{n-\frac{\alpha}{k}+1}}{\eta(\varphi(b), \varphi(a), m)} \int_0^1 t^{n-\frac{\alpha}{k}} |f^{(n+1)}(m\varphi(b) + t\eta(\varphi(x), \varphi(b), m))| dt \\ & \leq \frac{|\eta(\varphi(x), \varphi(a), m)|^{n-\frac{\alpha}{k}+1}}{\eta(\varphi(b), \varphi(a), m)} \left(\int_0^1 t^{(n-\frac{\alpha}{k})p} dt \right)^{\frac{1}{p}} \\ & \times \left(\int_0^1 |f^{(n+1)}(m\varphi(a) + t\eta(\varphi(x), \varphi(a), m))|^q dt \right)^{\frac{1}{q}} \\ & + \frac{|\eta(\varphi(x), \varphi(b), m)|^{n-\frac{\alpha}{k}+1}}{\eta(\varphi(b), \varphi(a), m)} \left(\int_0^1 t^{(n-\frac{\alpha}{k})p} dt \right)^{\frac{1}{p}} \\ & \times \left(\int_0^1 |f^{(n+1)}(m\varphi(b) + t\eta(\varphi(x), \varphi(b), m))|^q dt \right)^{\frac{1}{q}} \end{aligned}$$

$$\begin{aligned}
&\leq \frac{|\eta(\varphi(x), \varphi(a), m)|^{n-\frac{\alpha}{k}+1}}{\eta(\varphi(b), \varphi(a), m)} \left(\int_0^1 t^{(n-\frac{\alpha}{k})p} dt \right)^{\frac{1}{p}} \\
&\times \left(\int_0^1 \left[m(1-t)^s |f^{(n+1)}(\varphi(a))|^q + t^s |f^{(n+1)}(\varphi(x))|^q \right] dt \right)^{\frac{1}{q}} \\
&+ \frac{|\eta(\varphi(x), \varphi(b), m)|^{n-\frac{\alpha}{k}+1}}{\eta(\varphi(b), \varphi(a), m)} \left(\int_0^1 t^{(n-\frac{\alpha}{k})p} dt \right)^{\frac{1}{p}} \\
&\times \left(\int_0^1 \left[m(1-t)^s |f^{(n+1)}(\varphi(b))|^q + t^s |f^{(n+1)}(\varphi(x))|^q \right] dt \right)^{\frac{1}{q}} \\
&\leq \frac{M}{\left((n - \frac{\alpha}{k}) p + 1 \right)^{\frac{1}{p}}} \left(\frac{m+1}{s+1} \right)^{\frac{1}{q}} \\
&\times \left[\frac{|\eta(\varphi(x), \varphi(a), m)|^{n-\frac{\alpha}{k}+1} + |\eta(\varphi(x), \varphi(b), m)|^{n-\frac{\alpha}{k}+1}}{\eta(\varphi(b), \varphi(a), m)} \right].
\end{aligned}$$

□

Corollary 2.9. Under the same conditions as in Theorem 2.8, if we choose $m = k = 1$, $\eta(\varphi(y), \varphi(x), m) = \varphi(y) - m\varphi(x)$ and $\varphi(x) = x$, $\forall x \in I$, we get the following inequality for Caputo fractional derivatives:

$$\begin{aligned}
&\left| \left[\frac{(x-a)^{n-\alpha} - (x-b)^{n-\alpha}}{b-a} \right] f^{(n)}(x) \right. \\
&+ (-1)^{n+1} \frac{\Gamma(n-\alpha+1)}{b-a} \left. \left[{}^c D_{x-}^\alpha f(a) - {}^c D_{x-}^\alpha f(b) \right] \right| \\
(2.6) &\leq \frac{M}{\left((n - \alpha) p + 1 \right)^{\frac{1}{p}}} \left(\frac{2}{s+1} \right)^{\frac{1}{q}} \left[\frac{(x-a)^{n-\alpha+1} + (b-x)^{n-\alpha+1}}{b-a} \right].
\end{aligned}$$

Theorem 2.10. Let $\alpha > 0$, $k \geq 1$ and $\alpha \notin \{1, 2, 3, \dots\}$, $n = [\alpha] + 1$. Let $\varphi : I \rightarrow K$ be a continuous function. Suppose $K \subseteq \mathbf{R}$ be an open m -invex subset with respect to $\eta : K \times K \times (0, 1) \rightarrow \mathbf{R}$ for any fixed $s, m \in (0, 1]$. Assume that $f : K \rightarrow \mathbf{R}$ is a function on K° such that $f \in C^{n+1}[m\varphi(a), m\varphi(a) + \eta(\varphi(b), \varphi(a), m)]$, where $+\eta(\varphi(b), \varphi(a), m) > 0$.

If $|f^{(n+1)}|^q$ is generalized (s, m, φ) -preinvex function on $[m\varphi(a), m\varphi(a) + \eta(\varphi(b), \varphi(a), m)]$, $q \geq 1$ and $|f^{(n+1)}| \leq M$, then the following inequality for Caputo k -fractional derivatives holds:

$$\begin{aligned}
 & |I_{f,\eta,\varphi}(x; \alpha, k, n, m, a, b)| \\
 & \leq M \left(\frac{1}{n - \frac{\alpha}{k} + 1} \right)^{1-\frac{1}{q}} \left[m\beta \left(n - \frac{\alpha}{k} + 1, s + 1 \right) + \frac{1}{n + s - \frac{\alpha}{k} + 1} \right]^{\frac{1}{q}} \\
 (2.7) \quad & \times \left[\frac{|\eta(\varphi(x), \varphi(a), m)|^{n-\frac{\alpha}{k}+1} + |\eta(\varphi(x), \varphi(b), m)|^{n-\frac{\alpha}{k}+1}}{\eta(\varphi(b), \varphi(a), m)} \right].
 \end{aligned}$$

Proof. Suppose that $q \geq 1$. From Lemma 2.5, generalized (s, m, φ) -preinvexity of $|f^{(n+1)}|^q$, the well-known power mean inequality, upper bound of $|f^{(n+1)}|$ and properties of the modulus, we have

$$\begin{aligned}
 & |I_{f,\eta,\varphi}(x; \alpha, k, n, m, a, b)| \\
 & \leq \frac{|\eta(\varphi(x), \varphi(a), m)|^{n-\frac{\alpha}{k}+1}}{\eta(\varphi(b), \varphi(a), m)} \int_0^1 t^{n-\frac{\alpha}{k}} |f^{(n+1)}(m\varphi(a) + t\eta(\varphi(x), \varphi(a), m))| dt \\
 & + \frac{|\eta(\varphi(x), \varphi(b), m)|^{n-\frac{\alpha}{k}+1}}{\eta(\varphi(b), \varphi(a), m)} \int_0^1 t^{n-\frac{\alpha}{k}} |f^{(n+1)}(m\varphi(b) + t\eta(\varphi(x), \varphi(b), m))| dt \\
 & \leq \frac{|\eta(\varphi(x), \varphi(a), m)|^{n-\frac{\alpha}{k}+1}}{\eta(\varphi(b), \varphi(a), m)} \left(\int_0^1 t^{n-\frac{\alpha}{k}} dt \right)^{1-\frac{1}{q}} \\
 & \times \left(\int_0^1 t^{n-\frac{\alpha}{k}} |f^{(n+1)}(m\varphi(a) + t\eta(\varphi(x), \varphi(a), m))|^q dt \right)^{\frac{1}{q}} \\
 & + \frac{|\eta(\varphi(x), \varphi(b), m)|^{n-\frac{\alpha}{k}+1}}{\eta(\varphi(b), \varphi(a), m)} \left(\int_0^1 t^{n-\frac{\alpha}{k}} dt \right)^{1-\frac{1}{q}} \\
 & \times \left(\int_0^1 t^{n-\frac{\alpha}{k}} |f^{(n+1)}(m\varphi(b) + t\eta(\varphi(x), \varphi(b), m))|^q dt \right)^{\frac{1}{q}} \\
 & \leq \frac{|\eta(\varphi(x), \varphi(a), m)|^{n-\frac{\alpha}{k}+1}}{\eta(\varphi(b), \varphi(a), m)} \left(\int_0^1 t^{n-\frac{\alpha}{k}} dt \right)^{1-\frac{1}{q}}
 \end{aligned}$$

$$\begin{aligned}
& \times \left(\int_0^1 t^{n-\frac{\alpha}{k}} \left[m(1-t)^s \left| f^{(n+1)}(\varphi(a)) \right|^q + t^s \left| f^{(n+1)}(\varphi(x)) \right|^q \right] dt \right)^{\frac{1}{q}} \\
& + \frac{|\eta(\varphi(x), \varphi(b), m)|^{n-\frac{\alpha}{k}+1}}{\eta(\varphi(b), \varphi(a), m)} \left(\int_0^1 t^{n-\frac{\alpha}{k}} dt \right)^{1-\frac{1}{q}} \\
& \times \left(\int_0^1 t^{n-\frac{\alpha}{k}} \left[m(1-t)^s \left| f^{(n+1)}(\varphi(b)) \right|^q + t^s \left| f^{(n+1)}(\varphi(x)) \right|^q \right] dt \right)^{\frac{1}{q}} \\
& \leq M \left(\frac{1}{n - \frac{\alpha}{k} + 1} \right)^{1-\frac{1}{q}} \left[m\beta \left(n - \frac{\alpha}{k} + 1, s + 1 \right) + \frac{1}{n + s - \frac{\alpha}{k} + 1} \right]^{\frac{1}{q}} \\
& \times \left[\frac{|\eta(\varphi(x), \varphi(a), m)|^{n-\frac{\alpha}{k}+1} + |\eta(\varphi(x), \varphi(b), m)|^{n-\frac{\alpha}{k}+1}}{\eta(\varphi(b), \varphi(a), m)} \right].
\end{aligned}$$

□

Corollary 2.11. Under the same conditions as in Theorem 2.10, if we choose $m = k = 1$, $\eta(\varphi(y), \varphi(x), m) = \varphi(y) - m\varphi(x)$ and $\varphi(x) = x$, $\forall x \in I$, we get the following inequality for Caputo fractional derivatives:

$$\begin{aligned}
& \left| \left[\frac{(x-a)^{n-\alpha} - (x-b)^{n-\alpha}}{b-a} \right] f^{(n)}(x) \right. \\
& \left. + (-1)^{n+1} \frac{\Gamma(n-\alpha+1)}{b-a} \left[{}^c D_{x-}^\alpha f(a) - {}^c D_{x-}^\alpha f(b) \right] \right| \\
& \leq M \left(\frac{1}{n - \alpha + 1} \right)^{1-\frac{1}{q}} \left[\beta(n - \alpha + 1, s + 1) + \frac{1}{n + s - \alpha + 1} \right]^{\frac{1}{q}} \\
& \times \left[\frac{(x-a)^{n-\alpha+1} + (b-x)^{n-\alpha+1}}{b-a} \right]. \tag{2.8}
\end{aligned}$$

Corollary 2.12. Under the same conditions as in Theorem 2.10 for $q = 1$, we get Theorem 2.6.

3. Applications to special means

In the following we give certain generalizations of some notions for a positive valued function of a positive variable.

Definition 3.1. (see [43]) A function $M : \mathbf{R}_+^2 \rightarrow \mathbf{R}_+$, is called a Mean function if it has the following properties:

1. Homogeneity: $M(ax, ay) = aM(x, y)$, for all $a > 0$,
2. Symmetry: $M(x, y) = M(y, x)$,
3. Reflexivity: $M(x, x) = x$,
4. Monotonicity: If $x \leq x'$ and $y \leq y'$, then $M(x, y) \leq M(x', y')$,
5. Internality: $\min\{x, y\} \leq M(x, y) \leq \max\{x, y\}$.

We consider some means for arbitrary positive real numbers α, β ($\alpha \neq \beta$).

1. The arithmetic mean: $A := A(\alpha, \beta) = \frac{\alpha+\beta}{2}$
2. The geometric mean: $G := G(\alpha, \beta) = \sqrt{\alpha\beta}$
3. The harmonic mean: $H := H(\alpha, \beta) = \frac{2}{\frac{1}{\alpha} + \frac{1}{\beta}}$
4. The power mean: $P_r := P_r(\alpha, \beta) = \left(\frac{\alpha^r + \beta^r}{2}\right)^{\frac{1}{r}}, \quad r \geq 1$.
5. The identric mean: $I := I(\alpha, \beta) = \begin{cases} \frac{1}{e} \left(\frac{\beta^\beta}{\alpha^\alpha}\right), & \alpha \neq \beta; \\ \alpha, & \alpha = \beta. \end{cases}$
6. The logarithmic mean: $L := L(\alpha, \beta) = \frac{\beta - \alpha}{\ln(\beta) - \ln(\alpha)}$.
7. The generalized log-mean: $L_p := L_p(\alpha, \beta) = \left[\frac{\beta^{p+1} - \alpha^{p+1}}{(p+1)(\beta - \alpha)} \right]^{\frac{1}{p}}; \quad p \in \mathbf{R} \setminus \{-1, 0\}$.
8. The weighted p -power mean: $M_p \left(\begin{array}{cccc} \alpha_1, & \alpha_2, & \cdots, & \alpha_n \\ u_1, & u_2, & \cdots, & u_n \end{array} \right) = (\sum_{i=1}^n \alpha_i u_i^p)^{\frac{1}{p}}$
where $0 \leq \alpha_i \leq 1$, $u_i > 0$ ($i = 1, 2, \dots, n$) with $\sum_{i=1}^n \alpha_i = 1$.

It is well known that L_p is monotonic nondecreasing over $p \in \mathbf{R}$ with $L_{-1} := L$ and $L_0 := I$. In particular, we have the following inequality $H \leq G \leq L \leq I \leq A$. Now, let a and b be positive real numbers such that $a < b$. Consider the function $M := M(\varphi(x), \varphi(y)) : [\varphi(x), \varphi(x) + \eta(\varphi(y), \varphi(x))] \times [\varphi(x), \varphi(x) + \eta(\varphi(y), \varphi(x))] \rightarrow \mathbf{R}_+$, which is one of the above mentioned means and $\varphi : I \rightarrow K$ be a continuous function. Therefore one can obtain various inequalities using the results of Section 2 for these means as follows: Replace $\eta(\varphi(y), \varphi(x), m)$ with $\eta(\varphi(y), \varphi(x))$ and setting $\eta(\varphi(y), \varphi(x)) = M(\varphi(x), \varphi(y))$ for value $m = 1$ and $\forall x, y \in I$ in (2.5) and (2.7), one can obtain the following interesting inequalities involving means:

$$\begin{aligned}
(3.1) \quad & \left| I_{f,M(\cdot,\cdot),\varphi}(x; \alpha, k, n, 1, a, b) \right| \leq \frac{M}{\left((n - \frac{\alpha}{k}) p + 1 \right)^{\frac{1}{p}}} \left(\frac{2}{s+1} \right)^{\frac{1}{q}} \\
& \times \left[\frac{M^{n-\frac{\alpha}{k}+1}(\varphi(a), \varphi(x)) + M^{n-\frac{\alpha}{k}+1}(\varphi(b), \varphi(x))}{|M(\varphi(a), \varphi(b))|} \right], \\
(3.2) \quad & \left| I_{f,M(\cdot,\cdot),\varphi}(x; \alpha, k, n, 1, a, b) \right| \\
& \leq M \left(\frac{1}{n - \frac{\alpha}{k} + 1} \right)^{1-\frac{1}{q}} \left[\beta(n - \frac{\alpha}{k} + 1, s + 1) + \frac{1}{n + s - \frac{\alpha}{k} + 1} \right]^{\frac{1}{q}} \\
& \times \left[\frac{M^{n-\frac{\alpha}{k}+1}(\varphi(a), \varphi(x)) + M^{n-\frac{\alpha}{k}+1}(\varphi(b), \varphi(x))}{|M(\varphi(a), \varphi(b))|} \right].
\end{aligned}$$

Letting $M(\varphi(x), \varphi(y)) = A, G, H, P_r, I, L, L_p, M_p, \forall x, y \in I$, in (3.1) and (3.2), we get the inequalities involving means for a particular choice of a generalized $(s, 1, \varphi)$ -preinvex functions $|f^{(n+1)}|^q$. The details are left to the interested reader.

4. Conclusions

In the present paper, the notion of generalized (s, m, φ) -preinvex function was applied to established some new generalizations of Ostrowski type inequalities via Caputo k -fractional derivatives.

Motivated by this new interesting class of generalized (s, m, φ) -preinvex functions we can indeed see to be vital for fellow researchers and scientists working in the same domain.

We conclude that our methods considered here may be a stimulant for further investigations concerning Ostrowski and Hermite-Hadamard type integral inequalities for various kinds of preinvex functions involving classical integrals, Riemann-Liouville fractional integrals, k -fractional integrals, local fractional integrals, fractional integral operators, Caputo k -fractional derivatives, q -calculus, (p, q) -calculus, time scale calculus and conformable fractional integrals.

References

- [1] Ç. Yıldız, M. E. Özdemir and M. Z. Sarikaya, New generalizations of Ostrowski-like type inequalities for fractional integrals, *Kyungpook Math. J.*, **56**, pp. 161-172, (2016).
- [2] A. Kashuri and R. Liko, Ostrowski type fractional integral inequalities for generalized (s, m, φ) -preinvex functions, *Aust. J. Math. Anal. Appl.*, **13**, (1), Article 16, pp. 1-11, (2016).
- [3] A. Kashuri and R. Liko, Generalizations of Hermite-Hadamard and Ostrowski type inequalities for MT_m -preinvex functions, *Proyecciones*, **36**, (1), pp. 45-80, (2017).
- [4] M. Alomari, M. Darus, S. S. Dragomir and P. Cerone, Ostrowski type inequalities for functions whose derivatives are s -convex in the second sense, *Appl. Math. Lett.*, **23**, pp. 1071-1076, (2010).
- [5] R. P. Agarwal, M. J. Luo and R. K. Raina, On Ostrowski type inequalities, *Fasc. Math.*, **204**, pp. 5-27, (2016).
- [6] S. S. Dragomir, On the Ostrowski's integral inequality for mappings with bounded variation and applications, *Math. Ineq. & Appl.*, **1**, (2) (1998).
- [7] S. S. Dragomir, The Ostrowski integral inequality for Lipschitzian mappings and applications, *Comput. Math. Appl.*, **38**, pp. 33-37, (1999).

- [8] S. S. Dragomir, Ostrowski-type inequalities for Lebesgue integral: A survey of recent results, *Aust. J. Math. Anal. Appl.*, **14**, (1), pp. 1-287, (2017).
- [9] G. Farid, Some new Ostrowski type inequalities via fractional integrals, *Int. J. Anal. App.*, **14**, (1), pp. 64-68, (2017).
- [10] W. Liu, W. Wen and J. Park, Ostrowski type fractional integral inequalities for MT-convex functions, *Miskolc Math. Notes*, **16**, (1), pp. 249-256, (2015).
- [11] M. Matloka, Ostrowski type inequalities for functions whose derivatives are h -convex via fractional integrals, *Journal of Scientific Research and Reports*, **3**, (12), pp. 1633-1641, (2014).
- [12] A. A. Kilbas, H. M. Srivastava and J. J. Trujillo, Theory and applications of fractional differential equations, *North-Holland Math. Stud.*, **204**, Elsevier, New York-London, (2006).
- [13] G. Farid, A. Javed and A. U. Rehman, On Hadamard inequalities for ntimes differentiable functions which are relative convex via Caputo k -fractional derivatives, *Nonlinear Anal. Forum*, to appear.
- [14] Z. Liu, Some Ostrowski-Grüss type inequalities and applications, *Comput. Math. Appl.*, **53**, pp. 73-79, (2007).
- [15] M. E. Özdemir, H. Kavurmac and E. Set, Ostrowski's type inequalities for (α, m) -convex functions, *Kyungpook Math. J.*, **50**, pp. 371-378, (2010).
- [16] B. G. Pachpatte, On an inequality of Ostrowski type in three independent variables, *J. Math. Anal. Appl.*, **249**, pp. 583-591, (2000).
- [17] B. G. Pachpatte, On a new Ostrowski type inequality in two independent variables, *Tamkang J. Math.*, **32**, (1), pp. 45-49, (2001).
- [18] A. Rafiq, N. A. Mir and F. Ahmad, Weighted Čebyšev-Ostrowski type inequalities, *Applied Math. Mechanics (English Edition)*, **28**, (7), pp. 901-906, (2007).
- [19] N. Ujević, Sharp inequalities of Simpson type and Ostrowski type, *Comput. Math. Appl.*, **48**, pp. 145-151, (2004).

- [20] L. Zhongxue, On sharp inequalities of Simpson type and Ostrowski type in two independent variables, *Comput. Math. Appl.*, **56**, pp. 2043-2047, (2008).
- [21] T. Abdeljawad, On conformable fractional calculus, *J. Comput. Appl. Math.*, **279**, pp. 57-66, (2015).
- [22] Z. Dahmani, New inequalities in fractional integrals, *Int. J. Nonlinear Sci.*, **9**, (4), pp. 493-497, (2010).
- [23] Z. Dahmani, L. Tabharit and S. Taf, Some fractional integral inequalities, *Nonlinear Sci. Lett. A*, **1**, (2), pp. 155-160, (2010).
- [24] Z. Dahmani, On Minkowski and Hermite-Hadamard integral inequalities via fractional integration, *Ann. Funct. Anal.*, **1**, (1), pp. 51-58, (2010).
- [25] Z. Dahmani, L. Tabharit and S. Taf, New generalizations of Grüss inequality using Riemann-Liouville fractional integrals, *Bull. Math. Anal. Appl.*, **2**, (3), pp. 93-99, (2010).
- [26] U. N. Katugampola, A new approach to generalized fractional derivatives, *Bulletin Math. Anal. Appl.*, **6**, (4), pp. 1-15, (2014).
- [27] R. Khalil, M. Al Horani, A. Yousef and M. Sababheh, A new definition of fractional derivative, *J. Comput. Appl. Math.*, **264**, pp. 65-70, (2014).
- [28] S. D. Purohit and S. L. Kalla, Certain inequalities related to the Chebyshev's functional involving Erdelyi-Kober operators, *Scientia Series A: Math. Sci.*, **25**, pp. 53-63, (2014).
- [29] R. K. Raina, On generalized Wright's hypergeometric functions and fractional calculus operators, *East Asian Math. J.*, **21**, (2), pp. 191-203, (2005).
- [30] M. Z. Sarikaya, On the Ostrowski type integral inequality, *Acta Math. Univ. Comenianae*, **79**, (1), pp. 129-134, (2010).
- [31] E. Set, A. Gözpınar and J. Choi, Hermite-Hadamard type inequalities for twice differentiable m -convex functions via conformable fractional integrals, *Far East J. Math. Sci.*, **101**, (4), pp. 873-891, (2017).

- [32] T. S. Du, J. G. Liao and Y. J. Li, Properties and integral inequalities of Hadamard-Simpson type for the generalized (s, m) -preinvex functions, *J. Nonlinear Sci. Appl.*, **9**, pp. 3112-3126, (2016).
- [33] H. Hudzik and L. Maligranda, Some remarks on s -convex functions, *Aequationes Math.*, **48**, pp. 100-111, (1994).
- [34] M. Ahmadmir and R. Ullah, Some inequalities of Ostrowski and Grüss type for triple integrals on time scales, *Tamkang J. Math.*, **42**, (4), pp. 415-426, (2011).
- [35] S. S. Dragomir and S. Wang, A new inequality of Ostrowski's type in L_1 -norm and applications to some special means and to some numerical quadrature rules, *Tamkang J. Math.*, **28**, pp. 239-244, (1997).
- [36] S. S. Dragomir and S. Wang, An inequality of Ostrowski-Grüss type and its applications to the estimation of error bounds for some special means and for some numerical quadrature rules, *Comput. Math. Appl.*, **13**, (11), pp. 15-20, (1997).
- [37] Z. Liu, Some companions of an Ostrowski type inequality and applications, *J. Inequal. in Pure and Appl. Math.*, **10**, (2), Art. 52, 12 pp. ... (2009)
- [38] D. S. Mitrinovic, J. E. Pečarić and A. M. Fink, Classical and new inequalities in analysis, *Kluwer Academic Publishers, Dordrecht*, (1993).
- [39] T. Antczak, Mean value in invexity analysis, *Nonlinear Anal.*, **60**, pp. 1473-1484, (2005).
- [40] X. M. Yang, X. Q. Yang and K. L. Teo, Generalized invexity and generalized invariant monotonicity, *J. Optim. Theory Appl.*, **117**, pp. 607-625, (2003).
- [41] R. Pini, Invexity and generalized convexity, *Optimization*, **22**, pp. 513-525, (1991).
- [42] M. Tunç, Ostrowski type inequalities for functions whose derivatives are MT-convex, *J. Comput. Anal. Appl.*, **17**, (4), pp. 691-696, (2014).
- [43] P. S. Bullen, Handbook of Means and Their Inequalities, *Kluwer Academic Publishers, Dordrecht*, (2003).

Artion Kashuri

Department of Mathematics,
Faculty of Technical Science,
University "Ismail Qemali",
Vlora, *Albania*
e-mail : artionkashuri@gmail.com

and

Rozana Liko

Department of Mathematics,
Faculty of Technical Science,
University "Ismail Qemali",
Vlora, *Albania*
e-mail : rozanaliko86@gmail.com