

Proyecciones Journal of Mathematics
Vol. 37, N° 1, pp. 171-180, March 2018.
Universidad Católica del Norte
Antofagasta - Chile

Jordan triple derivation on alternative rings

Ruth N. Ferreira

Universidade Tecnológica Federal do Paraná, Brasil

and

Bruno L. M. Ferreira

Universidade Tecnológica Federal do Paraná, Brasil

Received : July 2017. Accepted : July 2017

Abstract

Let \mathbf{D} be a mapping from an alternative ring \mathbf{R} into itself satisfying $\mathbf{D}(a \cdot ba) = \mathbf{D}(a) \cdot ba + a \cdot \mathbf{D}(b)a + a \cdot b\mathbf{D}(a)$ for all $a, b \in \mathbf{R}$. Under some conditions on \mathbf{R} , we show that \mathbf{D} is additive.

AMS 2010 Subject Classification : *17A36, 17D05.*

Keywords : *Alternative ring; Idempotent element; Maps; Additivity.*

1. Introduction

In this paper, \mathbf{R} will be a ring not necessarily associative or commutative and consider the following convention for its multiplication operation: $xy \cdot z = (xy)z$ and $x \cdot yz = x(yz)$ for $x, y, z \in \mathbf{R}$, to reduce the number of parentheses. For $x, y, z \in \mathbf{R}$ we denote the *associator* by $(x, y, z) = (xy)z - x(yz)$.

A ring \mathbf{R} is called *k-torsion free* if $kx = 0$ implies $x = 0$, for any $x \in \mathbf{R}$, where $k \in \mathbf{Z}$, $k > 0$, *prime* if $IJ \neq 0$ for any two nonzero ideals $I, J \subseteq \mathbf{R}$ and *semiprime* if it contains no nonzero ideal whose square is zero.

A ring \mathbf{R} is said to be *alternative* if

$$(x, x, y) = 0 = (y, x, x), \text{ for all } x, y \in \mathbf{R},$$

and *flexible* if

$$(x, y, x) = 0, \text{ for all } x, y \in \mathbf{R}.$$

One easily sees that any alternative ring is flexible.

Theorem 1.1. *Let R be a 3-torsion free alternative ring. So R is a prime ring if and only if $aR \cdot b = 0$ (or $a \cdot Rb = 0$) implies $a = 0$ or $b = 0$ for $a, b \in R$.*

Proof. Clearly all alternative rings satisfying the properties $aR \cdot b = 0$ (or $a \cdot Rb = 0$) are prime rings. Suppose R is a prime ring by [7, Lemma 2.4, Theorem A and Proposition 3.5] we have $R = \mathbf{A}_0 \supseteq \mathbf{A}_1 \supseteq \cdots \supseteq \mathbf{A}_n = \mathbf{A} \neq (0)$ is a chain of subrings of R . If $aR \cdot b = 0$ (or $a \cdot Rb = 0$) hence $a\mathbf{A} \cdot b = 0$ (or $a \cdot \mathbf{A}b = 0$) follows [7, Proposition 3.5 (e)] that $a = 0$ or $b = 0$. \diamond

A mapping $\mathbf{D} : \mathbf{R} \rightarrow \mathbf{R}$ is Jordan triple multiplicative derivation if

$$\mathbf{D}(a \cdot ba) = \mathbf{D}(a) \cdot ba + a \cdot \mathbf{D}(b)a + a \cdot b\mathbf{D}(a)$$

for all $a, b \in \mathbf{R}$. It is worth noting that by the flexible identity of alternative rings we can write

$$\mathbf{D}(aba) = \mathbf{D}(a) \cdot ba + a\mathbf{D}(b)a + a \cdot b\mathbf{D}(a)$$

for all $a, b \in \mathbf{R}$. Let us consider \mathbf{R} an alternative ring and let us fix a nontrivial idempotent $e_1 \in \mathbf{R}$, i.e, $e_1^2 = e_1$; $e_1 \neq 0$ and e_1 is not an unity element. Let $e_2 : \mathbf{R} \rightarrow \mathbf{R}$ and $e'_2 : \mathbf{R} \rightarrow \mathbf{R}$ be given by $e_2a = a - e_1a$ and $e'_2a = a - ae_1$. We shall denote e'_2a by ae_2 . Note that \mathbf{R} need not have

an identity element. The operation $x(1 - y)$ for $x, y \in \mathbf{R}$ is understood as $x - xy$. It is easy to see that $(e_i a) e_j = e_i (a e_j)$ for all $a \in \mathcal{R}$ and $i, j = 1, 2$. Then \mathbf{R} has a Peirce decomposition $\mathbf{R} = \mathbf{R}_{11} \oplus \mathbf{R}_{12} \oplus \mathbf{R}_{21} \oplus \mathbf{R}_{22}$, where $\mathbf{R}_{ij} = e_i \mathbf{R} e_j$ ($i, j = 1, 2$), satisfying the multiplicative relations:

- (i) $\mathbf{R}_{ij} \mathbf{R}_{jl} \subseteq \mathbf{R}_{il}$ ($i, j, l = 1, 2$);
- (ii) $\mathbf{R}_{ij} \mathbf{R}_{ij} \subseteq \mathbf{R}_{ji}$ ($i, j = 1, 2; i \neq j$);
- (iii) $\mathbf{R}_{ij} \mathbf{R}_{kl} = 0$, if $j \neq k$ and $(i, j) \neq (k, l)$, ($i, j, k, l = 1, 2$);
- (iv) $x_{ij}^2 = 0$ for all $x_{ij} \in \mathbf{R}_{ij}$ ($i, j = 1, 2; i \neq j$).

Remark 1.1. By the linearization of (iv) we obtain

$$x_{ij} y_{ij} + y_{ij} x_{ij} = 0$$

if $i \neq j$. This identity is very useful for the main result to be verified.

The study of the relationship between the multiplicative and the additive structures of a ring has become an interesting and active topic in ring theory. In non-associative ring theory we can mention recent works such as [1], [2], [3], [4] where the authors generalize the results for a class of non-associative rings, namely alternative rings. The present paper we investigate the problem of when a Jordan triple multiplicative derivation must be an additive map for the class of alternative rings. The hypotheses of the main Theorem allow the author to make its proof based on calculus using the Peirce decomposition notion for Alternative rings. But it is worth noting that the notion of Peirce decomposition for the alternative rings is similar to the notion of Peirce decomposition for the associative rings. However, the similarity of this notion is only in its written form, but not in its theoretical structure because the Peirce decomposition for alternative rings is the generalization of the Peirce decomposition for associative rings. The symbol “.”, as defined in the introduction section of our article, is essential to elucidate how the non-associative multiplication should be done, and also the symbol “.” is used to simplify the notation. Therefore, the symbol “.” is crucial to the logic, characterization and generalization of associative results to the alternative results. In this paper we shall continue the line of research introduced in [5] and [6] where its authors demonstrate the following results

Theorem 1.2. Let \mathbf{R} be an alternative ring containing a non-trivial idempotent e_1 and $\mathbf{R} = \mathbf{R}_{11} \oplus \mathbf{R}_{12} \oplus \mathbf{R}_{21} \oplus \mathbf{R}_{22}$, the Peirce Decomposition of \mathbf{R} , relative to e_1 , satisfying:

(i) If $((e_ia)e_j)x_{jk} = 0$ for all $x_{jk} \in \mathbf{R}_{jk}$, then $((e_ia)e_j) = 0$;

(ii) If $x_{ij}((e_ja)e_i) = 0$ for all $x_{ij} \in \mathbf{R}_{ij}$, then $((e_ja)e_i) = 0$,

for $i, j, k \in \{1, 2\}$. If $D : \mathbf{R} \longrightarrow \mathbf{R}$ is a multiplicative derivation, then D is additive.

and

Theorem 1.3. Let \mathbf{R} be an alternative ring containing a non-trivial idempotent e_1 and $\mathbf{R} = \mathbf{R}_{11} \oplus \mathbf{R}_{12} \oplus \mathbf{R}_{21} \oplus \mathbf{R}_{22}$, the Peirce Decomposition of \mathbf{R} , relative to e_1 , satisfying:

(i) If $(e_iae_j)x_{jk} = 0$ for all $x_{jk} \in \mathbf{R}_{jk}$, then $(e_iae_j) = 0$;

(ii) If $x_{ij}(e_jae_k) = 0$ for all $x_{ij} \in \mathbf{R}_{ij}$, then $(e_jae_k) = 0$;

(iii) If $(e_iae_i)x_{ii} + x_{ii}(e_iae_i) = 0$ for all $x_{ii} \in \mathbf{R}_{ii}$, then $(e_iae_i) = 0$.

for $i, j, k \in \{1, 2\}$. If $\mathbf{D} : \mathbf{R} \longrightarrow \mathbf{R}$ is a Jordan multiplicative derivation, then \mathbf{D} is additive.

2. The main Theorem

We shall prove the following result:

Theorem 2.1. Let \mathbf{R} be an alternative ring containing a non-trivial idempotent e_1 and $\mathbf{R} = \mathbf{R}_{11} \oplus \mathbf{R}_{12} \oplus \mathbf{R}_{21} \oplus \mathbf{R}_{22}$, the Peirce Decomposition of \mathbf{R} , relative to e_1 , satisfying:

(i) If $(e_1ae_1)x_{12} = 0$ for all $x_{12} \in \mathbf{R}_{12}$, then $(e_1ae_1) = 0$;

(ii) If $x_{12}(e_2ae_2) = 0$ for all $x_{12} \in \mathbf{R}_{12}$, then $(e_2ae_2) = 0$;

(iii) Let $i, j \in \{1, 2\}$. If $x_{ij}ax_{ij} = 0$ for all $x_{ij} \in \mathbf{R}_{ij}$, then $(e_jae_i) = 0$.

If $\mathbf{D} : \mathbf{R} \longrightarrow \mathbf{R}$ is a Jordan triple multiplicative derivation, then \mathbf{D} is additive.

The proof of the Theorem is organized as a series of Lemmas.

We begin with the following Lemma with a simple proof.

Lemma 2.1. $\mathbf{D}(0) = 0$.

Proof. $D(0) = D(000) = \mathbf{D}(0) \cdot 00 + 0\mathbf{D}(0)0 + 0 \cdot 0\mathbf{D}(0) = 0.$ \diamond

Lemma 2.2. $\mathbf{D}(a_{11} + b_{12} + c_{21} + d_{22}) = \mathbf{D}(a_{11}) + \mathbf{D}(b_{12}) + \mathbf{D}(c_{21}) + \mathbf{D}(d_{22}).$

Proof. For any $x_{ij} \in \mathbf{R}$, $i, j = 1, 2$, on one hand, we have

$$\begin{aligned} \mathbf{D}[x_{ij}(a_{11} + b_{12} + c_{21} + d_{22})x_{ij}] &= \mathbf{D}(x_{ij}) \cdot (a_{11} + b_{12} + c_{21} + d_{22})x_{ij} \\ &+ x_{ij}\mathbf{D}(a_{11} + b_{12} + c_{21} + d_{22})x_{ij} \\ &+ x_{ij} \cdot (a_{11} + b_{12} + c_{21} + d_{22})\mathbf{D}(x_{ij}). \end{aligned}$$

On the other hand,

$$\begin{aligned} \mathbf{D}(x_{ij}a_{11}x_{ij}) &= \mathbf{D}(x_{ij}) \cdot a_{11}x_{ij} + x_{ij}\mathbf{D}(a_{11})x_{ij} + x_{ij} \cdot a_{11}\mathbf{D}(x_{ij}), \\ \mathbf{D}(x_{ij}b_{12}x_{ij}) &= \mathbf{D}(x_{ij}) \cdot b_{12}x_{ij} + x_{ij}\mathbf{D}(b_{12})x_{ij} + x_{ij} \cdot b_{12}\mathbf{D}(x_{ij}), \\ \mathbf{D}(x_{ij}c_{21}x_{ij}) &= \mathbf{D}(x_{ij}) \cdot c_{21}x_{ij} + x_{ij}\mathbf{D}(c_{21})x_{ij} + x_{ij} \cdot c_{21}\mathbf{D}(x_{ij}), \\ \mathbf{D}(x_{ij}d_{22}x_{ij}) &= \mathbf{D}(x_{ij}) \cdot d_{22}x_{ij} + x_{ij}\mathbf{D}(d_{22})x_{ij} + x_{ij} \cdot d_{22}\mathbf{D}(x_{ij}). \end{aligned}$$

These imply that

$$\begin{aligned} &\mathbf{D}[x_{ij}(a_{11} + b_{12} + c_{21} + d_{22})x_{ij}] - \mathbf{D}(x_{ij}a_{11}x_{ij}) \\ &- \mathbf{D}(x_{ij}b_{12}x_{ij}) - \mathbf{D}(x_{ij}c_{21}x_{ij}) - \mathbf{D}(x_{ij}d_{22}x_{ij}) \\ &= x_{ij}[\mathbf{D}(a_{11} + b_{12} + c_{21} + d_{22}) - \mathbf{D}(a_{11}) - \mathbf{D}(b_{12}) - \mathbf{D}(c_{21}) - \mathbf{D}(d_{22})]x_{ij}, \end{aligned}$$

where we use the flexible identity. By the flexible identity we note that for any $i, j = 1, 2$, we have

$$\begin{aligned} &\mathbf{D}[x_{ij}(a_{11} + b_{12} + c_{21} + d_{22})x_{ij}] - \mathbf{D}(x_{ij}a_{11}x_{ij}) \\ &- \mathbf{D}(x_{ij}b_{12}x_{ij}) - \mathbf{D}(x_{ij}c_{21}x_{ij}) - \mathbf{D}(x_{ij}d_{22}x_{ij}) = 0. \end{aligned}$$

Then, for $i, j = 1, 2$, we get

$$x_{ij}[\mathbf{D}(a_{11} + b_{12} + c_{21} + d_{22}) - \mathbf{D}(a_{11}) - \mathbf{D}(b_{12}) - \mathbf{D}(c_{21}) - \mathbf{D}(d_{22})]x_{ij} = 0.$$

By Condition (iii), we see that

$$[\mathbf{D}(a_{11} + b_{12} + c_{21} + d_{22}) - \mathbf{D}(a_{11}) - \mathbf{D}(b_{12}) - \mathbf{D}(c_{21}) - \mathbf{D}(d_{22})]_{ji} = 0, i, j = 1, 2.$$

Equivalently,

$$\mathbf{D}(a_{11} + b_{12} + c_{21} + d_{22}) - \mathbf{D}(a_{11}) - \mathbf{D}(b_{12}) - \mathbf{D}(c_{21}) - \mathbf{D}(d_{22}) = 0.$$

\diamond

- Lemma 2.3.** 1. $\mathbf{D}(a_{12} + b_{12}c_{22}) = \mathbf{D}(a_{12}) + \mathbf{D}(b_{12}c_{22})$,
 2. $\mathbf{D}(a_{21} + b_{22}c_{21}) = \mathbf{D}(a_{21}) + \mathbf{D}(b_{22}c_{21})$.

Proof. We will prove only (1) because the proof of (2) is similar to (1).
 By Remark 1.1 we note that

$$e_1 + a_{12} + b_{12}a_{12} + a_{12}b_{12} + b_{12}c_{22} = (e_1 + a_{12} + c_{22})(e_1 + b_{12})(e_1 + a_{12} + c_{22}).$$

By applying Lemma 2.2, Remark 1.1 and flexible identity we have

$$\begin{aligned} \mathbf{D}(e_1) + \mathbf{D}(a_{12} + b_{12}c_{22}) &= \mathbf{D}(e_1 + a_{12} + b_{12}c_{22}) \\ &= \mathbf{D}[(e_1 + a_{12} + c_{22})(e_1 + b_{12})(e_1 + a_{12} + c_{22})] \\ &= \mathbf{D}(e_1 + a_{12} + c_{22}) \cdot (e_1 + b_{12})(e_1 + a_{12} + c_{22}) \\ &+ (e_1 + a_{12} + c_{22})\mathbf{D}(e_1 + b_{12})(e_1 + a_{12} + c_{22}) \\ &+ (e_1 + a_{12} + c_{22}) \cdot (e_1 + b_{12})\mathbf{D}(e_1 + a_{12} + c_{22}) \\ &= [\mathbf{D}(e_1) + \mathbf{D}(a_{12}) + \mathbf{D}(c_{22})] \cdot (e_1 + b_{12})(e_1 + a_{12} + c_{22}) \\ &+ (e_1 + a_{12} + c_{22})[\mathbf{D}(e_1) + \mathbf{D}(b_{12})](e_1 + a_{12} + c_{22}) \\ &+ (e_1 + a_{12} + c_{22}) \cdot (e_1 + b_{12})[\mathbf{D}(e_1) + \mathbf{D}(a_{12}) + \mathbf{D}(c_{22})] \\ &= [\mathbf{D}(e_1) + \mathbf{D}(a_{12}) + \mathbf{D}(c_{22})] \cdot e_1(e_1 + a_{12} + c_{22}) \\ &+ (e_1 + a_{12} + c_{22})\mathbf{D}(e_1)(e_1 + a_{12} + c_{22}) \\ &+ (e_1 + a_{12} + c_{22}) \cdot e_1[\mathbf{D}(e_1) + \mathbf{D}(a_{12}) \\ &+ \mathbf{D}(c_{22})] + [\mathbf{D}(e_1) + \mathbf{D}(a_{12}) + \mathbf{D}(c_{22})] \cdot b_{12}(e_1 + a_{12} + c_{22}) \\ &+ (e_1 + a_{12} + c_{22})\mathbf{D}(b_{12})(e_1 + a_{12} + c_{22}) \\ &+ (e_1 + a_{12} + c_{22}) \cdot b_{12}[\mathbf{D}(e_1) + \mathbf{D}(a_{12}) + \mathbf{D}(c_{22})] \\ &= \mathbf{D}[(e_1 + a_{12} + c_{22})e_1(e_1 + a_{12} + c_{22})] \\ &+ \mathbf{D}[(e_1 + a_{12} + c_{22})b_{12}(e_1 + a_{12} + c_{22})] \\ &= \mathbf{D}(e_1 + a_{12}) + \mathbf{D}(b_{12}a_{12} + b_{12}c_{22} + a_{12}b_{12} + a_{12}b_{12}a_{12}) \\ &= \mathbf{D}(e_1) + \mathbf{D}(a_{12}) + \mathbf{D}(b_{12}c_{22}). \end{aligned}$$

To Prove (2) just use the identity

$$e_1 + a_{21} + b_{22}c_{21} = (e_1 + a_{21} + b_{22}) \cdot (e_1 + c_{21})(e_1 + a_{21} + b_{22}).$$

◇

Lemma 2.4. $\mathbf{D}(a_{ij} + b_{ij}) = \mathbf{D}(a_{ij}) + \mathbf{D}(b_{ij})$, $i \neq j$.

Proof. For any $x_{ij} \in \mathbf{R}_{ij}$, $i, j = 1, 2$, from

$$\mathbf{D}[x_{ij}(a_{ij} + b_{ij})x_{ij}] = 0 = \mathbf{D}(x_{ij}a_{ij}x_{ij}) + \mathbf{D}(x_{ij}b_{ij}x_{ij})$$

we can get

$$x_{ij}[\mathbf{D}(a_{ij} + b_{ij}) - \mathbf{D}(a_{ij}) - \mathbf{D}(b_{ij})]x_{ij} = 0.$$

This implies, by conditions of the Theorem 2.1 that

$$[\mathbf{D}(a_{ij} + b_{ij}) - \mathbf{D}(a_{ij}) - \mathbf{D}(b_{ij})]_{ii} = [\mathbf{D}(a_{ij} + b_{ij}) - \mathbf{D}(a_{ij}) - \mathbf{D}(b_{ij})]_{jj} = 0.$$

This completes the proof. \diamond

Lemma 2.5. $\mathbf{D}(a_{kk} + b_{kk}) = \mathbf{D}(a_{kk}) + \mathbf{D}(b_{kk})$.

Proof. Let a_{kk} and b_{kk} be arbitrary elements of \mathbf{R}_{kk} , $k = 1, 2$. By considering $\mathbf{D}[x_{ij}(a_{kk} + b_{kk})x_{ij}]$, $\mathbf{D}(x_{ij}a_{kk}x_{ij})$, and $\mathbf{D}(x_{ij}b_{kk}x_{ij})$ for the cases of $i \neq j$ and $i = j$ respectively, one can easily get that

$$[\mathbf{D}(a_{kk} + b_{kk}) - \mathbf{D}(a_{kk}) - \mathbf{D}(b_{kk})]_{ji} = 0,$$

$$[\mathbf{D}(a_{kk} + b_{kk}) - \mathbf{D}(a_{kk}) - \mathbf{D}(b_{kk})]_{jj} = 0,$$

where in the second identity $i = j \neq k$. Now we have only to prove that

$$[\mathbf{D}(a_{kk} + b_{kk}) - \mathbf{D}(a_{kk}) - \mathbf{D}(b_{kk})]_{jj} = 0,$$

with $k = j$. For any $x_{12} \in \mathbf{R}_{12}$, $r_{11} \in \mathbf{R}_{11}$ and $r_{22} \in \mathbf{R}_{22}$, from

$$(2.1) \quad r_{11}x_{12} = (e_1 + r_{11})x_{12}(e_1 + r_{11})$$

and

$$(2.2) \quad x_{12}r_{22} = (e_1 + r_{22})x_{12}(e_1 + r_{22})$$

can check, by (1) and (2) that

$$(2.3) \quad \begin{aligned} \mathbf{D}(r_{11}x_{12}) &= \mathbf{D}(e_1) \cdot r_{11}x_{12} + e_1 \cdot \mathbf{D}(r_{11})x_{12} + r_{11}\mathbf{D}(x_{12}) \\ &+ \mathbf{D}(x_{12})r_{11} + x_{12} \cdot \mathbf{D}(r_{11})e_1, \end{aligned}$$

$$(2.4) \quad \begin{aligned} \mathbf{D}(x_{12}r_{22}) &= \mathbf{D}(e_1) \cdot x_{12}r_{22} + e_1 \cdot \mathbf{D}(x_{12})r_{22} + r_{22} \\ &\cdot \mathbf{D}(x_{12})e_1 + x_{12}\mathbf{D}(r_{22}). \end{aligned}$$

Now, applying equality (3) for $r_{11} = a_{11} + b_{11}$, $r_{11} = a_{11}$ and $r_{11} = b_{11}$ and applying equality (4) for $r_{22} = a_{22} + b_{22}$, $r_{22} = a_{22}$ and $r_{22} = b_{22}$, we can get

$$\begin{aligned} [\mathbf{D}(a_{11} + b_{11}) - \mathbf{D}(a_{11}) - \mathbf{D}(b_{11})]x_{12} &= 0, \\ x_{12}[\mathbf{D}(a_{22} + b_{22}) - \mathbf{D}(a_{22}) - \mathbf{D}(b_{22})] &= 0. \end{aligned}$$

It follows from Condition (i) and (ii) of the Theorem 2.1 that $[\mathbf{D}(a_{kk} + b_{kk}) - \mathbf{D}(a_{kk}) - \mathbf{D}(b_{kk})]_{jj} = 0$, with $k = j$, which completes the proof. \diamond

Now we are ready to prove our main result.

Proof of the Theorem 2.1: For any $a, b \in \mathbf{R}$, we write $a = a_{11} + a_{12} + a_{21} + a_{22}$ and $b = b_{11} + b_{12} + b_{21} + b_{22}$. Applying the previous Lemmas, we have

$$\begin{aligned} \mathbf{D}(a + b) &= \mathbf{D}(a_{11} + a_{12} + a_{21} + a_{22} + b_{11} + b_{12} + b_{21} + b_{22}) \\ &= \mathbf{D}[(a_{11} + b_{11}) + (a_{12} + b_{12}) + (a_{21} + b_{21}) + (a_{22} + b_{22})] \\ &= \mathbf{D}(a_{11} + b_{11}) + \mathbf{D}(a_{12} + b_{12}) + \mathbf{D}(a_{21} + b_{21}) + \mathbf{D}(a_{22} + b_{22}) \\ &= \mathbf{D}(a_{11}) + \mathbf{D}(b_{11}) + \mathbf{D}(a_{12}) + \mathbf{D}(b_{12}) + \mathbf{D}(a_{21}) + \mathbf{D}(b_{21}) \\ &\quad + \mathbf{D}(a_{22}) + \mathbf{D}(b_{22}) \\ &= \mathbf{D}(a_{11} + a_{12} + a_{21} + a_{22}) + \mathbf{D}(b_{11} + b_{12} + b_{21} + b_{22}) \\ &= \mathbf{D}(a) + \mathbf{D}(b), \end{aligned}$$

i. e., \mathbf{D} is additive. \diamond

3. Applications in prime alternative rings

In the case of an unital alternative ring we have

Corollary 3.1. *Let \mathbf{R} be an unital alternative ring containing a non-trivial idempotent e_1 and $\mathbf{R} = \mathbf{R}_{11} \oplus \mathbf{R}_{12} \oplus \mathbf{R}_{21} \oplus \mathbf{R}_{22}$, the Peirce Decomposition of \mathbf{R} , relative to e_1 , satisfying:*

- (i) *If $(e_1ae_1)x_{12} = 0$ for all $x_{12} \in \mathbf{R}_{12}$, then $(e_1ae_1) = 0$;*
- (ii) *If $x_{12}(e_2ae_2) = 0$ for all $x_{12} \in \mathbf{R}_{12}$, then $(e_2ae_2) = 0$;*

(iii) Let $i, j \in \{1, 2\}$. If $x_{ij}ax_{ij} = 0$ for all $x_{ij} \in \mathbf{R}_{ij}$, ($i \neq j$) then $(e_jae_i) = 0$.

If $\mathbf{D} : \mathbf{R} \longrightarrow \mathbf{R}$ is a Jordan triple multiplicative derivation, then \mathbf{D} is additive.

As a last result of our paper follows the Corollary

Corollary 3.2. Let \mathbf{R} be a 3-torsion free prime unital alternative ring with a nontrivial idempotent. If mapping $\mathbf{D} : \mathbf{R} \rightarrow \mathbf{R}$ satisfies

$$\mathbf{D}(aba) = \mathbf{D}(a) \cdot ba + a\mathbf{D}(b)a + a \cdot b\mathbf{D}(a)$$

for all $a, b \in \mathbf{R}$, then \mathbf{D} is additive.

References

- [1] B. L. M. Ferreira and J. C. M. Ferreira, *Additivity of n -Multiplicative Maps on Alternative Rings*, Communications In Algebra **44**, pp. 1557-1568, (2016).
- [2] B. L. M. Ferreira, J. C. M. Ferreira and H. Guzzo Jr., *Jordan Maps on Alternative Algebras*, JP Journal of Algebra, Number Theory and Applications **31**, pp. 129-142, (2013).
- [3] B. L. M. Ferreira, J. C. M. Ferreira and H. Guzzo Jr., *Jordan Triple Elementary Maps on Alternative Rings*, Extracta Mathematicae **29**, pp. 1-18, (2014).
- [4] B. L. M. Ferreira, J. C. M. Ferreira and H. Guzzo Jr., *Jordan Triple Maps of Alternative Algebras*, JP Journal of Algebra, Number Theory and Applications **33**, pp. 25-33, (2014).
- [5] B. L. M. Ferreira and R. Nascimento, *Derivable Maps on Alternative Rings*, Recen **16**, pp. 9-15, (2014).
- [6] R. N. Ferreira and B. L. M. Ferreira, *Jordan Derivation on Alternative Rings*, International Journal of Mathematics, Game Theory and Algebra **25**, pp. 435-445, (2017).

- [7] M. Slater, *Prime Alternative Rings, I*, Journal of Algebra **15**, pp. 229-243, (1970).

Ruth N. Ferreira

Universidade Tecnológica Federal do Paraná,
Guarapuava,
Paraná
Brasil
e-mail : ruthnascimento@utfpr.edu.br

and

Bruno L. M. Ferreira

Universidade Tecnológica Federal do Paraná,
Guarapuava,
Paraná
Brasil
e-mail : brunoferreira@utfpr.edu.br